



Antarctic Treaty (Environment Protection) Proclamation 2005

Antarctic Treaty (Environment Protection) Act 1980

I, PHILIP MICHAEL JEFFERY, Governor-General of the Commonwealth of Australia, acting with the advice of the Federal Executive Council, make this Proclamation under subsection 8 (2) of the *Antarctic Treaty (Environment Protection) Act 1980*.

Signed and sealed with the
Great Seal of Australia
on 15 December 2005

P. M. JEFFERY
Governor-General

By His Excellency's Command

IAN MACDONALD
Minister for Fisheries, Forestry and Conservation
for the Minister for the Environment and Heritage

	Page
Contents	
1 Name of Proclamation	4
2 Commencement	4
3 Definition — <i>Act</i>	4
4 Declaration of Antarctic specially protected areas and Antarctic specially managed areas	4
5 Cessation of effect of certain declarations in Schedule 1	4
6 Revocation of certain earlier declarations	5
Schedule 1	
Descriptions of Antarctic specially protected areas	6
Part 1 Antarctic specially protected area No. 101	6
Part 2 Antarctic specially protected area No. 102	7
Part 3 Antarctic specially protected area No. 103	8
Part 5 Antarctic specially protected area No. 105	10
Part 6 Antarctic specially protected area No. 106	11
Part 7 Antarctic specially protected area No. 107	12
Part 8 Antarctic specially protected area No. 108	13
Part 9 Antarctic specially protected area No. 109	14
Part 10 Antarctic specially protected area No. 110	16
Part 11 Antarctic specially protected area No. 111	17
Part 12 Antarctic specially protected area No. 112	18
Part 13 Antarctic specially protected area No. 113	20
Part 14 Antarctic specially protected area No. 114	21
Part 15 Antarctic specially protected area No. 115	22
Part 16 Antarctic specially protected area No. 116	23
Part 17 Antarctic specially protected area No. 117	24
Part 18 Antarctic specially protected area No. 118	27
Part 19 Antarctic specially protected area No. 119	30
Part 20 Antarctic specially protected area No. 120	32
Part 21 Antarctic specially protected area No. 121	34
Part 22 Antarctic specially protected area No. 122	36
Part 23 Antarctic specially protected area No. 123	38
Part 24 Antarctic specially protected area No. 124	41
Part 25 Antarctic specially protected area No. 125	42
Part 26 Antarctic specially protected area No. 126	43
Part 27 Antarctic specially protected area No. 127	45
Part 28 Antarctic specially protected area No. 128	46
Part 29 Antarctic specially protected area No. 129	47
Part 30 Antarctic specially protected area No. 130	49
Part 31 Antarctic specially protected area No. 131	50
Part 32 Antarctic specially protected area No. 132	51
Part 33 Antarctic specially protected area No. 133	53
Part 34 Antarctic specially protected area No. 134	55
Part 35 Antarctic specially protected area No. 135	56

		Page
Part 36	Antarctic specially protected area No. 136	58
Part 37	Antarctic specially protected area No. 137	59
Part 38	Antarctic specially protected area No. 138	61
Part 39	Antarctic specially protected area No. 139	63
Part 40	Antarctic specially protected area No. 140	65
Part 41	Antarctic specially protected area No. 141	68
Part 42	Antarctic specially protected area No. 142	70
Part 43	Antarctic specially protected area No. 143	71
Part 44	Antarctic specially protected area No. 144	73
Part 45	Antarctic specially protected area No. 145	74
Part 46	Antarctic specially protected area No. 146	75
Part 47	Antarctic specially protected area No. 147	76
Part 48	Antarctic specially protected area No. 148	79
Part 49	Antarctic specially protected area No. 149	80
Part 50	Antarctic specially protected area No. 150	82
Part 51	Antarctic specially protected area No. 151	83
Part 52	Antarctic specially protected area No. 152	84
Part 53	Antarctic specially protected area No. 153	86
Part 54	Antarctic specially protected area No. 154	88
Part 55	Antarctic specially protected area No. 155	90
Part 56	Antarctic specially protected area No. 156	92
Part 57	Antarctic specially protected area No. 157	93
Part 58	Antarctic specially protected area No. 158	94
Part 59	Antarctic specially protected area No. 159	96
Part 60	Antarctic specially protected area No. 160	97
Part 61	Antarctic specially protected area No. 161	98
Part 63	Antarctic Specially Protected Area No. 163	100
Part 64	Antarctic Specially Protected Area No. 164	101
Schedule 2	Antarctic specially managed areas	105
Part 2	Antarctic Specially Managed Area No. 2	105
Part 3	Antarctic Specially Managed Area No. 3	108
Part 4	Antarctic Specially Managed Area No. 4	110

1 Name of Proclamation

This Proclamation is the *Antarctic Treaty (Environment Protection) Proclamation 2005*.

2 Commencement

This Proclamation commences on 1 January 2006.

3 Definition — Act

In this Proclamation:

Act means the *Antarctic Treaty (Environment Protection) Act 1980*.

4 Declaration of Antarctic specially protected areas and Antarctic specially managed areas

- (1) Each area described in Schedule 1, being an area in respect of which the Antarctic Treaty Consultative Parties have adopted a management plan, is declared to be an Antarctic specially protected area.
- (2) Each area described in Schedule 2, being an area in respect of which the Antarctic Treaty Consultative Parties have adopted a management plan, is declared to be an Antarctic specially managed area.
- (3) For paragraph 8 (4) (a) of the Act, each Antarctic specially protected area and Antarctic specially managed area includes the subsoil beneath the surface of any land within the area, to a depth of 1 000 metres below the surface of the land.
- (4) For paragraph 8 (4) (c) of the Act, each Antarctic specially protected area and Antarctic specially managed area includes the subsoil beneath any sea-bed within the area, to a depth of 1 000 metres below the sea-bed.

Note Complete details of Antarctic specially protected areas and Antarctic specially managed areas (including management plans) are available at <http://davey.nettas.net/apa/index.html>.

5 Cessation of effect of certain declarations in Schedule 1

The declarations, as Antarctic specially protected areas, of the areas described in Parts 25, 27, 44, 46 and 50 of Schedule 1 cease to have effect at the end of 31 December 2010.

6 Revocation of certain earlier declarations

- (1) The declarations made by the *Antarctic Treaty (Environment Protection) Proclamation 2002* and the *Antarctic Treaty (Environment Protection) Amendment Proclamation 2004 (No. 1)* are revoked.

Note The Proclamations were published in the *Gazette* as follows:

- the *Antarctic Treaty (Environment Protection) Proclamation 2002* — on 8 January 2003;
- the *Antarctic Treaty (Environment Protection) Amendment Proclamation 2004 (No. 1)* — on 23 December 2004.

- (2) The declarations (other than the declaration, as an Antarctic specially protected area, of the area described, in the Proclamation made under the Act on 4 December 1980, as Antarctic specially protected area No. 4) taken, under subsection 8 (5) of the Act, to have been made by a Proclamation or Proclamations made on 24 November 2002 are revoked.

Note 1 Section 11 of the *Antarctic Treaty (Environment Protection) Legislation Amendment Act 1992*, which substituted a new section 8 into the Act, commenced on 24 November 2002.

Note 2 That Proclamation was published in the *Gazette* on 20 January 1981 (note that the text of the Proclamation published in the *Gazette* incorrectly shows the date of the making of the Proclamation as 4 December 1981).

Schedule 1 Descriptions of Antarctic specially protected areas

(section 4)

Part 1 Antarctic specially protected area No. 101

1 Name and location


Taylor Rookery, MacRobertson Land, East Antarctica.

Latitude 67° 27' S, longitude 60° 52' E.

2 Description of values to be protected

Taylor Rookery contains the largest Emperor penguin colony known on land; almost all other emperor penguins colonies are located on sea ice. The number of breeding pairs at the colony has ranged from 2 462 in 1989 to 3 307 in 1990 and has averaged approximately 3 000 over 15 years from 1988 to 2002. The rookery is important because of long-term monitoring of the population of the penguins (since 1954). The colony is ideal for counting since it is surrounded by small rocky hills which make it possible to observe every bird without entering the breeding area. A photographic census program has been carried out annually since 1988, and it is believed that this method has resulted in almost complete accuracy of counting.

3 Map



Part 2 Antarctic specially protected area No. 102

1 Name and location

Rookery Islands, MacRobertson Land, East Antarctica.

Latitude 67° 37' S, longitude 62° 32' E.

2 Description of values to be protected


The Rookery Islands contain breeding colonies of six bird species resident in the Mawson area: Adélie penguin (*Pygoscelis adeliae*), Cape petrel (*Daption capense*), snow petrel (*Pagodroma nivea*), southern giant petrel (*Macronectes giganteus*), Wilson's storm petrel (*Oceanites oceanicus*) and the Antarctic skua (*Catharacta maccormicki*). The southern giant petrel breeds nowhere else in the region. The designation of the Area aims to safeguard this unusual association of six species and ensure the preservation of a representative offshore island habitat.

The Rookery Islands provides a representative sample of the offshore island habitats occurring along the coast of MacRobertson land.

The southern giant petrel (*Macronectes giganteus*) has a world population of approximately 62,000 individuals and is inferred to have sustained a population reduction of at least 20% over the last 60 years. The species is in continued rapid decline. Giganteus Island in the Rookery Islands group is one of only four known breeding localities of southern giant petrels around the coastline of continental Antarctica. The other three continental breeding colonies are located near the Australian stations of Casey (66° 13' S, 110° 11' E), (Frazier Islands) and Davis (68° 35' S, 77° 58' E) (Hawker Island), and near the French station Dumont d'Urville (66° 40' S, 140° 01' E) in Terre Adélie. The current population for continental Antarctica is estimated at approximately 290 pairs, comprised of 3 pairs on Giganteus Island, 25 pairs on Hawker Island, 16 pairs at Pointe Géologie archipelago (Terre Adélie) and 248 pairs on the Frazier Islands. Southern giant petrels on the Antarctic continent comprise less than 1% of the global breeding population.

Southern giant petrels are widespread in more northerly latitudes, breeding on islands to the north-west of the Antarctic Peninsula and on islands of the Scotia Ridge. However, it is important that it should be protected at the southern limit of its breeding range.

3 Map



Part 3 Antarctic specially protected area No. 103

1 Name and location

Ardery Island and Odbert Island, Budd Coast, Wilkes Land, East Antarctica.


Latitude 66° 22' S, longitude 110° 27' E.

2 Description of values to be protected

Ardery Island and Odbert Island support several breeding species of petrel. There is no other readily accessible place in eastern Antarctica where the four genera of fulmarine petrels (*Thalassoica antarctica*, *Fulmarus glacialis*, *Daption capense* and *Pagodroma nivea*) breed in the same place in sufficient numbers to allow comparative study. Study of these four genera at one location is of high ecological importance in understanding and monitoring the Southern Ocean ecosystem.

It is believed that Ardery Island is unique insofar as it is the only area in the Antarctic which harbours two different subspecies of snow petrels. Studies on morphological or ecological differences between these two subspecies are not possible anywhere else. In addition both islands have breeding populations of Wilson's storm petrels (*Oceanites oceanicus*) and Antarctic skuas (*Catharacta maccormicki*) and Odber Island supports breeding populations of Adélie penguins (*Pygoscelis adeliae*).

3 Map



Note Part 4 is reserved.

Part 5 Antarctic specially protected area No. 105

1 Name and location


Beaufort Island, Ross Sea.

Latitude 76° 59' S, longitude 167° 00' E.

2 Description

Beaufort Island measures 7 km by 3.2 km. The designated Area encompasses the whole of Beaufort Island (76° 59' S, 167° 00' E) above the mean high water mark, and includes adjacent fast ice occupied by breeding Emperor penguins.

3 Map



Part 6 Antarctic specially protected area No. 106

1 Name and location


Cape Hallett, Victoria Land.

Latitude 72° 19' S, longitude 170° 16' E.

2 Description

Cape Hallett is located at the southern end of Moubray Bay, Northern Victoria Land, in the western Ross Sea. The Area includes Seabee Hook and the adjacent western slopes up to the crest of the ridge of the north end of the Hallett Peninsula, east of Willett Cove to the margin of the permanent ice sheet. The sea-level boundary is predominantly defined by the coastline of Seabee Hook, and extends south along the east shore of Willett Cove to the southern-most boundary at 72° 19' 30" S (approximately 800 m south of the Moubray Bay coast). The east boundary of the Area follows the edge of the permanent ice sheet near the ridge of Hallett Peninsula.

3 Map



Note This map predates the current terminology — Specially Protected Area No. 7 is now Antarctic specially protected area No. 106.

Part 7 Antarctic specially protected area No. 107

1 Name and location

Dion Islands, Marguerite Bay, Antarctic Peninsula.


Latitude 67° 53' S, longitude 68° 42' W.

2 Description

The Dion Islands (latitude 67° 53' S, longitude 68° 42' W, within a region of approximately 12 km²), situated 13.5 km south of the south-western extremity of Adelaide Island in north-western Marguerite Bay, is a small archipelago comprising the following islands, rocks and reefs: Envoy Rock, Regent Reef, Consort Islands (approximately 3 ha), Emperor Island (approximately 5 ha), Jester Rock, Noble Rocks, Courtier Islands (approximately 8 ha), Embassy Islands and Consul Reef.

The designated Area of just over 3 km² comprises Emperor Island and the marine environment (including sea ice when present) within 1 000 m of the coastline of Emperor Island. However, the area does not include the terrestrial areas of the Consort Islands in the north, Jester Rock in the east, or the Courtier Islands in the southwest.

3 Map



Part 8 Antarctic specially protected area No. 108

1 Name and location

Green Island, Berthelot Islands, Antarctic Peninsula.


Latitude 65° 19' S, longitude 64° 09' W.

2 Description

Green Island (65° 19' S, 64° 09' W, approximately 0.2 km²) is a small island situated 150 m north of the largest of the Berthelot Islands group, Grandidier Channel, approximately 3 km off the Graham Coast of the Antarctic Peninsula.

The designated area comprises the entire island, with the boundary defined as the low tide level. Offshore islets and rocks are not included within the Area. Boundary markers have not been installed. The coast itself is a clearly defined and visually obvious boundary feature.

3 Map



Part 9 Antarctic specially protected area No. 109

1 Name and location


Moe Island, South Orkney Islands.

Latitude 60° 44' S, longitude 45° 41' W.

2 Description

Moe Island, South Orkney Islands, is a small irregularly-shaped island lying 300 m off the south-western extremity of Signy Island, from which it is separated by Fyr Channel. It is about 1.3 km from the north-east to south-west and 1 km from north-west to south-east. Its position on Admiralty Chart No. 1775, latitude 60° 44' S, longitude 45° 45' W, does not agree closely with that in the map (latitude 60° 44' S, longitude 45° 41' W).

3 Map



Note This map predates the current terminology — Specially Protected Area No. 13 is now Antarctic specially protected area No. 109.

Part 10 Antarctic specially protected area No. 110

1 Name and location

Lynch Island, South Orkney Islands.


Latitude 60° 39' 10" S, longitude 45° 36' 25" W.

2 Description

Lynch Island (latitude 60° 39' 10" S, longitude 45° 36' 25" W; area: 0.1 km²) is a small island situated at the eastern end of Marshall Bay in the South Orkney Islands, about 200 m south of Coronation Island and 2.4 km north of Signy Island.

The designated Area comprises the entire island above the low tide level, at which the coastline is defined as the boundary of the Area. Boundary markers have not been installed because the coast itself is a clearly defined and visually obvious boundary feature.

3 Map



Note This map predates the current terminology — SPA is now ASPA.

Part 11 Antarctic specially protected area No. 111

1 Name and location


Southern Powell Island and adjacent islands, South Orkney Islands.

Latitude 60° 42' S, longitude 45° 01' W.

2 Description

The Area, which is centred on latitude 60° 42' S and longitude 45° 01' W, includes all of Powell Island, South Orkney Islands, south of the latitude of the southern summit of John Peaks (375 m altitude), together with the shore of Fredriksen Island, Michelsen Island (a tidal peninsula at the southern tip of Powell Island), Christoffersen Island, Grey Island and unnamed adjacent islands.

3 Map



Note This map predates the current terminology — SPA No. 15 is now ASPA No. 111.

Part 12 Antarctic specially protected area No. 112

1 Name and location


Coppermine Peninsula, Robert Island, South Shetland Islands.

Latitude 62° 23' S, longitude 59° 42' W.

2 Description

Coppermine Peninsula ($62^{\circ} 23' \text{ S}$, $59^{\circ} 42' \text{ W}$) is situated on the west side of Robert Island, which lies between Nelson Island to the east and Greenwich Island to the west, midway along the South Shetland Islands archipelago. The Area comprises all land west of a north-south line across the isthmus between Carlota Cove and Coppermine Cove, 100 m west of a small group of Chilean refuge huts. The Peninsula is about 1.7 km from south-east to north-west and up to 0.6 km from north-east to south-west, and is largely surrounded by precipitous cliffs.

3 Map



Note This map predates the current terminology — SPA No. 16 is now ASPA No. 112.

Part 13 Antarctic specially protected area No. 113


1 Name and location

Litchfield Island, Arthur Harbour, Anvers Island, Palmer Archipelago.

Latitude 64° 46' S, longitude 64° 06' W.

2 Description

Approximate area: 2.7 km². A small island designated on the grounds that, together with its littoral zone, it possesses an unusually rich collection of marine and terrestrial life, is unique amongst the neighbouring islands as a breeding place for six species of native birds, and provides an outstanding example of the natural ecological system of the Antarctic Peninsula area.



Note ASPA No. 113 was formerly specially protected area No. 17 — see the Proclamation of 4 December 1980.

Part 14 Antarctic specially protected area No. 114

1 Name and location


Northern Coronation Island, South Orkney Islands.

Latitude 60° 33' S, longitude 45° 35' W.

2 Description

The Area includes the region of northern Coronation Island between Conception Point to the west and Foul Point to the east. The eastern boundary follows a ridge from Foul Point approximately 6 km southwards to the summit of Mount Nivea (1 266 m), thence west-south-westwards for a distance of 1 500 m down the ridge to the col at High Stile. From High Stile, the boundary continues west-south-westwards for approximately 6 km following the ridge of the broad plateau of Brisbane Heights to the summit of Wave Peak (960 m). From Wave Peak the boundary extends due north for 1 000 m, thence west and in a north-westerly direction for about 6 km following the broad ridge of Brisbane Heights. The boundary then extends due north for approximately 6 km, following the main ridgeline to Conception Point. The glacial catchments draining to the northern coastline of Coronation Island within this boundary are within the Area. The actual summits of Mount Nivea and Wave Peak and the southern side of High Stile are outside of the Area. The northern boundary is defined as a straight line extending 11 km across the sea from Conception Point to Foul Point, including Ommanney Bay and the bay further to the west as within the Area.

3 Map



Part 15 Antarctic specially protected area No. 115

1 Name and location

Lagotellerie Island, Marguerite Bay, Antarctic Peninsula.


Latitude 67° 53' 20" S, longitude 67° 25' 30" W.

2 Description

Lagotellerie Island (latitude 67° 53' 20" S, longitude 67° 25' 30" W; area 1.58 km²), is situated in Marguerite Bay, Fallières Coast, Graham Land, 46 km south-east of Rothera Point on Adelaide Island, 11 km south of Porquois Pas Island and 3.25 km west of the south end of Horseshoe Island. Lagotellerie Island is 2 km by 1.3 km, oriented generally in an east-west direction.

The designated Area comprises the entire main island, and offshore islets within 200 m of the coast, above the low tide water level, which is defined as the boundary of the Area. Boundary markers have not been installed because the coast itself is a clearly defined and visually obvious boundary.

3 Map



Part 16 Antarctic specially protected area No. 116

1 Name and location


New College Valley, Caughley Beach, Cape Bird, Ross Island.

Latitude 77° 10' S, longitude 166° 40' E.

2 Description

Caughley Beach and its hinterland lie between the areas known as the Cape Bird Northern and Middle Penguin Rookeries, about 1 km north of Cape Bird, northern Ross Island (latitude 77° 10' S, longitude 166° 40' E). The proposed site encompasses the area between the top of the coastal cliffs of Caughley Beach and the Mt Bird Ice Cap, and between a line 200 m south of the New Zealand Antarctic Research Programme's summer station and a line 500 m north of Cape Bird Middle Adélie Penguin Rookery.

3 Map



Note This map predates the current terminology — SPA No. 20 is now ASPA No. 116.

Part 17 Antarctic specially protected area No. 117

1 Name and location

Avian Island, off Adelaide Island, Antarctic Peninsula.


Latitude 67° 46' S, longitude 68° 54' W.

2 Description

Avian Island (latitude 67° 46' S, longitude 68° 54' W, 0.49 km²), is situated in the north-west of Marguerite Bay, 400 m south of the south-western extremity of Adelaide Island. The island is 1.45 km long by 0.8 km at its widest, and is of roughly triangular shape.

The designated Area comprises the whole of Avian Island and the littoral zone, offshore islets and rocks, and a buffer zone of the surrounding marine environment (including sea ice when present) within 100 m of the shoreline of the main island. Boundary markers have not been installed because the coast forms a visually obvious reference for the marine boundary.

3 Map



Part 18 Antarctic specially protected area No. 118

1 Name and location


Summit of Mount Melbourne, Victoria Land.


Latitude 74° 21' S, longitude 164° 42' E.

2 Description

Mount Melbourne (2 733 m, 74° 21' S, 164° 42' E) in northern Victoria Land, is situated between Wood Bay and Terra Nova Bay, on the western side of the Ross Sea, and Campbell Glacier, about 10 km to the west. The Area encompasses all terrain above the 2 200 m contour surrounding the main crater of Mt Melbourne.

3 Maps





Note These maps do not appear in the management plan for APSA No. 118. They are included here for reference.

Part 19 Antarctic specially protected area No. 119

1 Name and location

Davis Valley and Forlidas Pond, Dufek Massif.

Latitude 82°27' S, longitude 51°21' W.

2 Description of values to be protected

The Davis Valley and the adjacent ice-free valleys is one of the most southerly 'dry valley' systems in Antarctica and, as of May 2005, is the most southerly protected area in Antarctica. While occupying an area of only 53 km², which is less than 1% of the area of the McMurdo Dry Valleys, the Area nevertheless contains the largest ice-free valley system found south of 80° S in the 90° W–0°–90° E half of Antarctica. Moreover, it is the only area known in this part of Antarctica where the geomorphology preserves such a detailed record of past glacial history. Some ice-free areas around the Weddell Sea region have scattered erratics and sometimes moraines, but the assemblage of drift limits, moraines, and abundant quartz-bearing erratics in the Davis Valley and associated valleys is unique and rare. The location of the Dufek Massif close to the junction between the western and the eastern Antarctic ice sheets also makes this site particularly valuable for the collection of data that can be used to constrain parameters such as the past thickness and dynamics of this sector of the Antarctic ice sheet. Such data are potentially extremely valuable for understanding the response of the Antarctic ice sheet to climate change. The Area therefore has exceptional and unique scientific value for the interpretation of past glacial events and climate in this part of Antarctica and it is important that this value is maintained.

The terrestrial ecology of the Area is impoverished but is also highly unusual, with lake and meltwater stream environments and their associated biota being rare this far south in Antarctica. As such, they provide unique opportunities for the scientific study of biological communities near the extreme limit of the occurrence of these environments. Vegetation appears to be limited to cyanobacterial mats and a very sparse occurrence of small crustose lichens. The cyanobacterial mat growth in the terrestrial locations is surprisingly extensive, and represents the best examples of this community type known this far south. The cyanobacterial community appears to survive in at least three distinct environments:

- in the permanent water bodies;
- in exposed terrestrial locations, particularly at the boundaries of sorted polygons; and
- in a series of former or seasonally dry pond beds on ice-free ground in the Davis Valley.

No arthropods or nematodes have thus far been detected in samples taken from within the Area, and the invertebrate fauna in the Area is unusually sparse. This characteristic distinguishes the Area from more northerly ice-free valley systems such as those at the Ablation Valley — Ganymede Heights (ASPA No. 147), Alexander Island, or at the McMurdo Dry Valleys (ASMA No. 2), where such communities are present. Rotifers and tardigrades have been extracted from samples taken within the Area, with the greatest numbers occurring within the former pond beds in the Davis Valley, although their diversity and abundance is also extremely limited compared with more northerly Antarctic sites (Hodgson and Convey, 2004). Further analyses of the samples obtained and identification of all taxa present are in progress, and are expected to make an important contribution to the understanding of biogeographical relationships between the different regions of Antarctica.


The Area is extremely isolated and difficult to access, and as a result has been visited by only a small number of people. Reports indicate that small field parties visited the Area in December 1957, in the 1965–66 and 1973–74 austral summer seasons, in December 1978 and in December 2003. The total number of people having visited probably numbers less than 50, with visits generally limited to a period of a few weeks or days. No structures or installations have been built within the Area, and as far as is known all equipment brought into the Area has subsequently been removed. While Hodgson and Convey (2004) reported evidence of a very limited number of human footprints and several old soil pit excavations, the Area has been exposed to few opportunities for direct human impact. The Area is believed to be one of the most pristine ice-free valley systems in Antarctica, and is therefore considered to possess outstanding potential as a reference area for microbiological studies, and it is important that these values receive long-term protection.

The site possesses outstanding wilderness and aesthetic values. The dry and weathered brown valleys of the Area are surrounded by extensive ice-fields, the margins of which fringe the valleys with dry based glacial ice of a deep blue hue. This abrupt and dramatic blue-ice margin stands in stark contrast to the stony and barren ice-free landscape of the valleys, and aesthetically is extremely striking in appearance. One of the original explorers of this area in 1957 recalled “the excitement we felt at being the first people to view and enter this magnificently scenic, pristine area.” (Behrendt, 1998: 354). Further examples of descriptions of the Area by visitors are: “[the blue ice] was towering over us — 150 feet — a large wave of blue. It was like being in a tidal wave that was held in suspension as we walked under it...” (Reynolds, field notes, 1978), and “I still cannot find adequate superlatives to describe the features, whether large or small, biologic or physical... [Of the] many settings that stretch the imagination...in my experience none match the northern side of the Dufek Massif, with Davis Valley as its crown jewel.” (Reynolds, pers. comm., 2000); “the most unusual [landscape] I have ever seen on any of the seven continents.” (Boyer, pers. comm., 2000); “Probably the single most remarkable environment I’ve been, either in Antarctica or elsewhere” (Convey, pers.

comm., 2004). Burt (2004) described the region simply as “inspiringly awesome”.

The boundaries of the Area have been revised to include the entire ice-free region centered on the Davis Valley, including the adjacent valleys and Forlidas Pond. In general, the margins of the surrounding ice sheets form the new boundary of the Area, resulting in special protection of the region as an integrated ice-free unit that more closely approximates the valley catchments. The full catchments of the surrounding glaciers that flow into these valleys extend considerable distances from the ice-free area and do not possess many of the values related to the purpose of special protection, and are therefore excluded from the Area.

3 Map



Part 20 Antarctic specially protected area No. 120

1 Name and location

Pointe-Geologie, Terre Adelie.

Latitude 66° 40' S, longitude 140° 02' E.

2 Description of values to be protected

In 1995, four islands, a nunatak and a breeding ground for emperor penguins were classified as an Antarctic Specially Protected Area (Measure 3 (1995), XIX ATCM, Seoul) because they were a representative example of terrestrial Antarctic ecosystems from a biological, geological and aesthetics perspective. A species of marine mammal, the Weddell seal (*Leptonychotes weddelli*) and various species of birds breed in the area: emperor penguin (*Aptenodytes forsteri*); Antarctic skua (*Catharacta maccormicki*); Adélie penguins (*Pygoscelis adeliae*); Wilson's petrel (*Oceanites oceanicus*); giant petrel (*Macronectes giganteus*); snow petrel (*Pagodroma nivea*); cape petrel (*Daption capense*).


Well-marked hills display asymmetrical transverse profiles with gently dipping northern slopes compared to the steeper southern ones. The terrain is affected by numerous cracks and fractures leading to very rough surfaces. The basement rocks consist mainly of sillimanite, cordierite and garnet-rich gneisses which are intruded by abundant dikes of pink anatexites. The lowest parts of the islands are covered by morainic boulders with a heterogenous granulometry (from a few centimetres to more than a metre across).

Long-term research and monitoring programs of birds and marine mammals have been going on for a long time already (since 1952 or 1964 according to the species). A database implemented in 1981 is directed by the Centre d'Etudes Biologiques de Chize (CEBC-CNRS). Human scientific presence may be estimated at around 4 people for a few hours, twice a month in the protected area.

Among the approximately 30 emperor penguin breeding sites on record, this is the only one located adjacent to a permanent station. It is therefore a providential spot to study this species and its environment.

3 Map

The Cape Geology Archipelago map shows the boundaries of the Antarctic Specially Protected Area inside the archipelago with dotted lines.



Part 21 Antarctic specially protected area No. 121

1 Name and location

Cape Royds, Ross Island.

Latitude 77° 33' 20" S, longitude 166° 09' 56" E.

2 Description


Cape Royds (77° 33' 20" S, 166° 09' 56" E) is situated on the west side of Ross Island, McMurdo Sound, at the western extremity of a coastal strip of ice-free land approximately 8 km wide, on the west slope of Mount Erebus. The Area comprises both a terrestrial and marine component.

The marine component of the Area extends for approximately 5 km from Derrick Point in the south to Rocky Point in the north, including Horseshoe Bay. The marine boundary is defined as extending north-east along the Arrival Bay coastline from the most easterly corner of the terrestrial boundary at Arrival Bay (77° 33' 15.9" S, 166° 10' 06" E) to Derrick Point (77° 33' 14.1" S, 166° 10' 22" E). From Derrick Point the marine boundary extends offshore 500 m in a south-east direction and thence extends parallel to the coast 500 m offshore from the mean high water mark, around Cape Royds and north for 5.3 km to a point 500 m due north of Rocky Point then due south to Rocky Point.

The terrestrial component of the Area consists of ice-free land within approximately 350 m of Cape Royds itself (77° 33' 20" S, 166° 09' 56" E). Much of this land is seasonally occupied by a breeding Adélie penguin colony. The boundary of this part of the Area has been revised from the original description so as to include all of the area occupied by breeding penguins in 1995/96 and the main southern access route of the penguins to the sea. The north boundary of the territorial component of the Area extends 45 m from a small embayment, on the west side and 350 m north of Cape Royds, in a straight line north-east to a survey mark identified on earlier New Zealand maps as IT2 (77° 33' 11.1" S, 166° 09' 33.3" E), which is an iron tube embedded in the ground. This line extends 10 m east from IT2 to a signpost (77° 33' 11.1" S, 166° 09' 34.8" E), then a further 80 m east to a signpost (77° 33' 11.0" S, 166° 09' 46.1" E) at the south end of a small pond at the north end of Pony Lake. From this signpost the boundary extends in a south-east direction for 114 m just north of the lake to the eastern edge of the lake (77° 33' 12.6" S, 166° 10' 01.3" E). The east boundary then extends 86 m in a south-south-east direction to a third signpost (77° 33' 15.2" S, 166° 10' 05" E), thence to the coast on the east side of Arrival Bay (77° 33' 15.9" S, 166° 10' 06.0" E). All of the ice-free ground, snow-patches and freshwater bodies contained west and south of the line defined above to the coast extending around Cape Royds is included within the Area.

3

Map



Part 22 Antarctic specially protected area No. 122

1


Name

Arrival Heights, Hut Point Peninsula, Ross Island.


2 Description

All that area of Arrival Heights enclosed with a line drawn from Trig T510 north-west over First crater to the 500-foot contour, then north along this contour to a point immediately west of Second Crater, then around the lip of this crater and south to Trig T510. The boundary, which is demarcated, is shown on the maps.

3 Maps



Note This map predates the current terminology — SSSI No. 2 is now ASPA No. 122.



Part 23 Antarctic specially protected area No. 123


- 1 **Name**
Barwick Valley, Victoria Land.

2 Description

Barwick Valley is situated about 65 km inland from the Ross Sea coast of South Victoria Land. The Area includes Barwick and Balham Valleys and their respective catchments and is bordered on the south, west and north by the McKelvey Valley, the Willet Range and the divide between the Victoria and Barwick Valleys, respectively.

The boundary of the Area extends from its eastern extremity in the lower Barwick Valley (around the confluence of the Barwick, Victoria and McKelvey Valleys) several kilometres south towards the ridge leading south-west to the summit of Mount Insel (1 345 m), from where the boundary follows the high points of the ridge of the Insel Range for 5 km before descending to a low pass between the McKelvey and Balham Valleys at the location of Bullseye Lake. The boundary crosses the lake before ascending the ridge to a further high point on the Insel Range (approximately 1 250 m), and continues towards the upper reaches of the Balham Valley. As the terrain becomes gentler in the upper Balham, the boundary sweeps north approximately above the 1 800 m contour line. The boundary skirts around the summit of Shapeless Mountain until it cuts north-west at a point west of the Apocalypse Peaks. The boundary joins and follows a prominent ridge to the summit of Mount Bastion (2 477 m, 160° 34' E, 77° 19' S). This ridge is followed in a northerly direction to Skew Peak (2 535 m, 160° 26' E, 77° 13' S), located at the head of the Barwick Valley. The boundary then descends along the East Ridge of Skew Peak above Webb Glacier, before following the catchment boundary in a more southerly direction toward Parker Mesa. From Parker Mesa the boundary descends further to follow the dividing ridge between the catchments of the Victoria Upper Glacier and the Barwick Valley. The boundary extends east along this ridge for 13 km to Sponsors Peak (1 454 m, 161° 24' E, 77° 18' S). The boundary descends the south-west Ridge of Sponsors Peak and Nickell Peak (approximately 1 400 m) to the lower Barwick to the eastern extremity of the Area, which is about 3 km north-east of Lake Vida, Victoria Valley.

3 Map



Note This map predates the current terminology — Site of Special Scientific Interest No. 3 is now Antarctic specially protected area No. 123.

Part 24 Antarctic specially protected area No. 124

1 **Name**


Cape Crozier, Ross Island.

2 **Description**

Cape Crozier is at the eastern extremity of Ross Island, where an ice-free area comprises the lower eastern slopes of Mount Terror. The designated area is situated in the vicinity of Post Office Hill (407 m), extending to encompass the adjacent Ross Ice Shelf where large cracks in the shelf are covered by fast ice which is occupied annually by breeding emperor penguins.

The Area includes a terrestrial region and ice shelf above the mean high water mark as well as the adjacent fast ice within the boundaries occupied by breeding emperor penguins. The north boundary of the Area extends 6.5 km along the 77° 26' 03" S line of latitude from 169° 11' 43" E to 169° 28' 00" E. The west boundary extends 1.5 km south from the northern boundary to the coast, thence in a north-east direction following a low ice-free ridge that passes 30 m west of the hut and helicopter pad. The boundary then follows this ridge in a southerly direction to be south-west of the summit of Post Office Hill before following another ridge in a south-east direction to the summit of a small unnamed peak (335 m) 1.2 km south-south-east of Post Office Hill. The boundary descends down a south-west ridgeline, before following this ridge to ascend to the summit of a large unnamed volcanic cone (520 m) 3.6 km south of Post Office Hill. The boundary follows around the eastern side of this cone before descending south through a valley to another unnamed conic peak (580 m) 1.5 km north-north-east of Bomb Peak. The boundary follows a ridge through the middle of the cone before descending down a ridge on the southern side of the peak into a valley at the base of Bomb Peak. The boundary ascends the northern side of Bomb Peak to the summit (>610 m) before extending down a ridge line on the south-east side of Bomb Peak to Igloo Spur and thence due east along latitude 77° 32' 00" S to the east boundary at 169° 28' 00" E.

3 Map



Part 25 Antarctic specially protected area No. 125

Note The declaration of this area as an Antarctic specially protected area ceases to have effect at the end of 31 December 2010 — see section 5.


1 Name

Fildes Peninsula, King George Island, South Shetland Islands.

2 Description

The two areas on Fildes Peninsula shown on the map will be demarcated.

3 Map



Note This map predates the current terminology — SSSI No. 5 is now ASPA No. 125.

Part 26 Antarctic specially protected area No. 126

1 Name and location

Byers Peninsula, Livingston Island, South Shetland Islands.


Latitude 62° 34' 35" S to 62° 40' 35" S, longitude 60° 54' 14" W to 61° 13' 07" W.

2 Description

Byers Peninsula (between latitudes 62° 34' 35" and 62° 40' 35" S and longitudes 60° 54' 14" and 61° 13' 07" W, 60.6 km²) is situated at the west end of Livingston Island, the second largest of the South Shetland Islands. The peninsula has a central west-east extent of about 9 km and a north-west south-east extent of 18.2 km, and is the largest ice-free area in the South Shetland Islands.

The boundaries of the Area designated under Recommendation XVI-5 have been changed. The Area now includes two islets several hundred metres south-west of Devils Point and a small area of ice-free ground at Clark Nunatak in the south-east corner as these sites also support values consistent with the remainder of the Peninsula. The Area is now defined to include the whole of Byers Peninsula west of the permanent ice sheet of Rotch Dome, Livingston Island, above the low tide water level, including the two islets adjacent to Devils Point noted above, but excluding all other offshore islets and rocks.

3 Map



Part 27 Antarctic specially protected area No. 127

Note The declaration of this area as an Antarctic specially protected area ceases to have effect at the end of 31 December 2010 — see section 5.

1 Name and location


Haswell Island.

Latitude 66° 31' S, longitude 93° 00' E.

2 Description

The site consists of Haswell Island (66° 31' S, 93° 00' E), about 1 km² in area, the largest of a group of islands lying close to Mirny station, together with its littoral zone and the area of fast ice, when present, lying within the delimitation shown on the map.

3 Map



Note This map predates the current terminology — SSSI No. 7 is now ASPA No. 127.

Part 28 Antarctic specially protected area No. 128

1 Name

Western shore of Admiralty Bay, King George Island.

2 Description

The area consists of land on the western shore of Admiralty Bay. The westerly boundary extends from Patelnia (Telefon) Point (62° 13' 55" S, 58° 28' 45" W), north-north-west to The Tower (a distinctive peak above Tower Glacier, 366.9 m at 62° 12' 50" S, 58° 29' 00" W), then continuing in a straight line to encompass the base of Jardine Peak (62° 10' 05" S, 58° 29' 45" W). This line then runs north-east to the sea (Admiralty Bay) where it bisects the coast immediately north of Rakusa Point (62° 09' 45" S, 58° 27' 25" W). Thereafter the Area is all the land which is bounded by the coastline south towards Demay Point (62° 12' 50" S, 58° 25' 15" W), then south-west along the coast to Patelnia (Telefon) Point. The western edge of the Area is adjacent to the Warsaw Icefield, with the north-western corner being ice-free in the vicinity of Jardine Peak. Outside of the northern boundary is a small area of ice-free land. Steep cliffs overlook a narrow beach and the waters of Ezcurra Inlet in the north-western section of the ice-free area; and in the north-eastern section there are occasional shallow beaches which extend to the sea, where H. Arctowski station is located, 400 m outside of the Area. Three small glaciers, Ecology, Baranowski, and Tower, descend from the Warsaw Icefield onto these shores. There are markers on the northern edge of the Area where the site has a boundary on land, immediately south of H. Arctowski station. The western boundary is not delineated by virtue of fact that it traverses a high (ca. 350 m) mobile icefield. The coastline defines the Area's eastern and southern stretches.

3 Map



Part 29 Antarctic specially protected area No. 129

1 Name and location


Rothera Point, Adelaide Island.

Latitude 67° 34' S, longitude 68° 08' W.

2 Description

Rothera Point (latitude 67° 34' S, longitude 68° 08' W) is situated in Ryder Bay, at the south-east corner of Wright Peninsula on the east side of Adelaide Island, south-west Antarctic Peninsula. The site is the north-eastern one-third of Rothera Point, and is representative of the area as a whole. It is about 30 m from west to east and 250 m from north to south, and rises to a maximum height of 36 m. At the coast, the site boundary is the 2.5 m contour. No upper shore, littoral or sublittoral areas of Rothera Point are therefore included. The southern boundary of the site, running across Rothera Point, is marked by a line of pink fuel drums filled with concrete. The remaining boundary is unmarked.

3 Map



Note This map predates the current terminology — Site of Special Scientific Interest No. 9 is now Antarctic specially protected area No. 129 and SSSI is now ASPA.

Part 30 Antarctic specially protected area No. 130

1 Name and location


Tramway Ridge, Mount Erebus, Ross Island.

Latitude 77° 31' 05" S, longitude 167° 06' 35" E.

2 Description

The boundary of the designated Area is defined as a square of 200 m by 200.8 m which encompasses most of the warm ground area of lower Tramway Ridge (77° 31' 05" S, 167° 06' 35" E). The Area is divided into two parts of almost equal size, the northern half being a Prohibited Zone. The boundaries of the Area and the Prohibited Zone (marked by signposts at each corner) and prominent features are shown on the map. Several boundary signposts have been offset owing to dangerous ground at the actual corner point.

3 Map



Part 31 Antarctic specially protected area No. 131

1 Name and location


Canada Glacier, Lake Fryxell, Taylor Valley, Victoria Land.

Latitude 77° 37' S, longitude 163° 03' E.

2 Description

The designated Area encompasses most of the glacier forefield area on the east side of the lower Canada Glacier, on the north shore of Lake Fryxell (77° 37' S, 163° 03' E). The south boundary of the Area is defined as the shoreline of Lake Fryxell, to the water's edge, extending about 1 km east from where the Canada Glacier toe meets Lake Fryxell. The west boundary of the Area follows the edge of the Canada Glacier. The south-east corner is near the neck of a small peninsula extending into Lake Fryxell. The peninsula, outside of the Area, is marked by a large rock (split) surrounded by a circle of rocks which was a benchmark for the 1985 NZ survey of the original SSSI. A wooden post marking Dry Valley Drilling Project Site 7 (1973) is about 10 m to the north-west of this point. A moraine ridge extends from the south-east corner upward and in a northerly direction: this ridge defines the eastern boundary of the Area. The ridge dips sharply before joining the featureless slope of the main Taylor Valley wall: the north-east corner of the Area is in this dip and will be marked by a cairn. A cairn will also be placed on a knoll on the ridge 450 m from the south-east corner point. Above the central flush a slope of lateral moraine of fairly even gradient extends upward and parallel to the glacier for about 1 km. At the top of this slope is a small knoll (220 m) about 300 m from where the glacier emerges into the Taylor Valley: a cairn and signpost will mark the boundary of the Area at this point. The upper, northern boundary of the Area extends from the Canada Glacier to the boundary markers on the knoll and thence declines in elevation in an easterly direction for 1.7 km to the north-east boundary cairn.

3 Map



Note This map predates the current terminology — Site of Special Scientific Interest No. 12 (SSSI No. 12) is now Antarctic specially protected area No. 131.

Part 32 Antarctic specially protected area No. 132

1 Name and location

Potter Peninsula, King George Island, South Shetland Islands.

Latitude 62°15' S, longitude 59°39' W.

2 **Description of values to be protected**

The area encompasses important scientific values due to the presence of groups of breeding elephant seals (*Mirounga leonina*), and non-breeding groups of Antarctic fur seals (*Arctocephalus gazella*) and occasionally of Weddell seals (*Leptonychotes weddelli*), crabeater seals (*Lobodon carcinophagus*) and leopard seals (*Hydrurga leptonyx*). During the breeding season there are around 500 elephant seals around, and during the fledging season, between 200 and 600. The non-breeding groups of Antarctic fur seals can add up to 300 individuals, although that figure may vary considerably from one year to the next.

Also present are significant colonies of gentoo penguins (*Pygoscelis papua*) and Adélie penguins (*Pygoscelis adeliae*), each of them with an approximate population of 4 000 pairs. Penguin colonies are mainly distributed around Stranger Point, from the vicinity of Elefante refuge to Spur 7. Within the ASPA, the population of storm petrels (by large *Oceanites oceanicus* and, to a much lesser extent, the *Fregetta tropica*) reaches some 200 pairs. Antarctic terns (*Sterna vittata*), southern giant petrels (*Macronectes giganteus*) and skuas (*Catharacta* sp.) also breed in the area, although their populations are, in all cases, less than 100 pairs each. Southern giant petrel nesting sites concentrate on the western side of the ASPA. As some of the nesting sites around Potter Peninsula change their position over time, population figures are estimations and may introduce some degree of inaccuracy, depending on the year considered.

3 Map

Map 1: General location of Antarctic Specially Protected Area No. 132, Potter Peninsula (King George Island), in relation to the Antarctic Peninsula.



Part 33 Antarctic specially protected area No. 133

1 Name and location

Harmony Point, Nelson Island, South Shetland Islands.

Latitude 62°18' S, longitude 59°11' W.

2 Description of values to be protected

Geomorphologically, Harmony Point presents three well defined units: an andesitic plateau, coastal and shelf outcrops and ancient sea levels. The plateau reaches 40 metres above sea level and its area is covered by detritus resulting from the action of erosion agents on andesitic rocks, with a well-developed vegetation of mosses and lichens.

There are three successive raised paleobeaches, between the coast and the glacier. The paleobeaches are defined by pebble accumulations of variable height in some instances and the development of soil in others.

Lakes and streams with a limited flow appear on the undulations. Some isolated andesitic rocks and ancient nunataks can be seen outside the limits of the glacier, evidencing that the past extension of the glacier covered Harmony Point.


The Area holds breeding colonies of 12 species: 3347 pairs of gentoo penguins (*Pygoscelis papua*), 89 685 pairs of chinstrap penguins (*Pygoscelis antarctica*), 746 pairs of southern giant petrels (*Macronectes giganteus*), 479 pairs of cape petrel (*Daption capense*), 45 pairs of blue eyed shag (*Phalacrocorax atriceps*), 144 pairs of snowy sheathbill (*Chionis alba*), 71 pairs of skuas (61 pairs of Antarctic skuas (*Catharacta antarctica*) and 11 of Polar skuas), 128 pairs of Dominican gull (*Larus dominicanus*) and between 100 and 150 individuals of Antarctic terns (*Sterna vittata*).

Other seabirds nesting in the Area are the Wilson's storm petrel (*Oceanites oceanicus*) and the black-bellied storm petrel (*Fregetta tropica*). Together they represent around 1 000 pairs. Map 4 shows the distribution of bird concentrations at Harmony Point.

There are usually 3 species of mammals in the Area: the Weddell seal (*Leptonychotes weddelli*), the elephant seal (*Mirounga leonina*) and the Antarctic fur seal (*Arctocephalus gazella*). Occasionally some individual crabeater seals (*Lobodon carcinophagus*) have also been seen. The number of mammals in the area varies. The maximum numbers of Antarctic fur seals, Weddell seals and elephant seals are 320, 550 and 100 respectively. The Weddell seals usually breed in the Area in high figures, reaching up to 60 females with their pups in a single season. Births of fur and elephant seals have also been recorded, although the numbers there are much lower.

There are some extensive areas covered by a very rich and diverse development of bryophytes and lichen-dominated plant communities (presently being classified), including, although to a lesser degree, two vascular plant species (*Deschampsia antarctica* and *Colobanthus quitensis*), especially in the areas less affected by recent anthropic perturbation or breeding activities. Moss turf subformations are located in wind protected and moist places, whilst lichen-dominated subformations occur in places with a high wind exposure. Five soil orders have been identified so far in the Area, according to the taxonomic system: Soil Taxonomy (1999): histosols (hydric cryfibrists), entisols (lithic criorthents), spodosols (oxiaquic humicryods), mollisols (lithic haplocryolls) and inceptisols (lithic eutrocrypts e histic cryaquepts).

3 Map



Part 34 Antarctic specially protected area No. 134

1 Name and location


Cierva Point and offshore islands, Danco Coast, Antarctic Peninsula.

Latitude 64° 10' S, longitude 60° 57' W (Cierva Point).

2 Description

Cierva Point (latitude 64° 10' S, longitude 60° 57' W) is in the north-west of the Antarctic Peninsula, on the south coast of Cierva Cove, at the north end of Hughes Bay. The site comprises Cierva Point, encompassing the land west of an imaginary line drawn from the south-east of the north side of the point, through the summit of a flat hill 540 m high, to the south-east of the south side of the point. Also included are Apéndice/Rivera/Sterneck Island, and José Hernández/Bofill/Midas Island and Cierva Point. Although the intertidal zone of each of these areas is included in the Area, the subtidal marine environment is not included. Base Primavera (Argentina) and all its associated installations and areas of disturbance are excluded from the Area.

3 Map



Part 35 Antarctic specially protected area No. 135

1 Name and location

North-east Bailey Peninsula, Budd Coast, Wilkes Land.


Latitude 66° 17' S, longitude 110° 32' E.

2 Description

The Area consists of an irregular area of exposed rock during summer on the north-east of Bailey Peninsula, with the north-western portion of the Area approximately 70 metres south of Brown Bay with Casey Station (66° 16' 59.9" S, 110° 31' 59.9" E) approximately 200 metres to the west. Boundary coordinates for the Area are shown in the following table.

Boundary point	Longitude	Latitude
1	110° 32' 42" E	66° 17' 03" S
2	110° 32' 56" E	66° 17' 11" S
3	110° 32' 50" E	66° 17' 11" S
4	110° 32' 41" E	66° 17' 10" S
5	110° 32' 22" E	66° 17' 07" S
6	110° 32' 20" E	66° 17' 06" S
7	110° 32' 18" E	66° 17' 02" S
8	110° 32' 18" E	66° 17' 00" S
9	110° 32' 14" E	66° 16' 60" S
10	110° 32' 09" E	66° 16' 56" S
11	110° 32' 08" E	66° 16' 54" S
12	110° 32' 05" E	66° 16' 54" S
13	110° 32' 07" E	66° 16' 52" S
14	110° 32' 07" E	66° 16' 52" S
15	110° 32' 12" E	66° 16' 51" S
16	110° 32' 16" E	66° 16' 52" S
17	110° 32' 19" E	66° 16' 53" S
18	110° 32' 19" E	66° 16' 55" S
19	110° 32' 24" E	66° 16' 55" S
20	110° 32' 25" E	66° 16' 53" S
21	110° 32' 29" E	66° 16' 53" S
22	110° 32' 44" E	66° 16' 54" S
23	110° 33' 09" E	66° 17' 05" S
24	110° 33' 11" E	66° 17' 06" S
25	110° 33' 10" E	66° 17' 09" S
26	110° 33' 02" E	66° 17' 11" S
27	110° 32' 56" E	66° 17' 11" S

3 Map



Part 36 Antarctic specially protected area No. 136

1 Name and location


Clark Peninsula, Budd Coast, Wilkes Land.

Latitude 66° 15' S, longitude 110° 36' E.

2 Description

Clark Peninsula, an area of rock exposures and permanent ice and snow fields, is situated on the north side of Newcomb Bay at the east end of Vincennes Bay, opposite Windmill Islands region, on Budd Coast, Wilkes Land, in latitude 66° 15' S and longitude 110° 36' E. The Area is approximately 9.75 km² in area. The Area comprises all the land on Clark Peninsula within the southern boundary line connecting the east side of Powell Cove at a point which originates at latitude 66.254424° S, longitude 110.53330° E, to trigonometrical station G7 at latitude 66.25809° S, longitude 110.55664° E thence to a point to the east-south-east on Løken Moraine. The eastern boundary is the western-most limit of Løken Moraines as far north as a point due east of Blakeney Point, and thence to the coast, returning along the coast to the point of origin. The boundary of the Area will be indicated by prominent markers.

3 Map



Note This map predates the current terminology — Site of Special Scientific Interest No. 17 is now Antarctic specially protected area No. 136.

Part 37 Antarctic specially protected area No. 137

1 Name and location

North-west White Island, McMurdo Sound.

Latitude 78° 00' S, longitude 167° 20' E.


2 Description

White Island, part of the McMurdo volcanic complex, is situated approximately 20 km south-east of the edge of the McMurdo Ice Shelf and 25 km south-east of Hut Point, the location of McMurdo Station (United States) and Scott Base (New Zealand) on Ross Island. The roughly triangular island is approximately 30 km long and 15 km wide at its maximum, and rises to a maximum elevation of 762 m in several locations.

The island is predominantly ice-covered with most of the rock outcrops being in the north, and is completely surrounded by the permanent shelf ice, between 10 m and 100 m in thickness, of the McMurdo Ice Shelf and Ross Ice Shelf. Black Island is situated 2.5 km west of White Island, separated by the shelf ice of White Strait.

The Area includes 130 km² of the shelf ice and open-water cracks of both the Ross Ice Shelf and McMurdo Ice Shelf up to 5 km offshore north-east, north and west from the White Island coast. The Area extends along the coast from a northern point on the east side of Cape Spencer-Smith (78° 00' 43" S, 167° 32' 42" E) 19 km south to the southern-most significant coastal outcrop of rock on the north-west side of White Island (78° 09' 08" S, 167° 05' 00" E). At this southern point, the boundary extends approximately 1 km due west to 167° 00' 00" E longitude, from where it extends due north to 78° 05' 00" S latitude, after which the boundary continues north at a constant distance of 5 km from the shore of White Island to the eastern extremity of the Area at 77° 58' 48" S, 167° 41' 35" E. The White Island coast is distinguished by a change in surface slope where the transition between the floating ice-shelf and land occurs: the transition is in some places gradual and indistinct, and the exact position of the coast is not precisely known. For this reason the coastal (generally east) boundary of the Area is considered to follow the line of the coast as evidenced by a surface elevation rise towards the land of two metres above the average elevation of the adjacent McMurdo Ice Shelf.

3



Part 38

1

Linnaeus Terrace, Asgaard Range, Victoria Land.


Latitude 77° 35' 50" S, longitude 161° 05' 00"E.

2 Description

Linnaeus Terrace (77° 35' 50" S, 161° 05' 00" E) is an elevated bench of weathered Beacon Sandstone approximately 1.5 km in length and 1 km in width. It is located at the east end of the Asgaard Range, 1.5 km north of Oliver Peak (77° 36' 40" S, 161° 02' 30" E) at an elevation of about 1 600 m. The Area overlooks the South Fork of the Wright Valley, is about 4 km from Don Juan Pond and 10 km from the terminus of the Wright Upper Glacier.

On the ground, the lower (northern) boundary of the Area is characterised by the presence of a predominantly sandstone outcrop of approximately 3 m in height which extends for much of the length of the terrace. The lower boundary of the Area is defined as the upper edge of this outcrop, and as straight lines adjoining the visible edges where the outcrop is covered by surface talus. The upper (southern) boundary of the Area is characterised by a line of sandstone outcrop of about 2–5 m in height, occurring between the elevations of 1 660–1 700 m about 70 m above the general elevation of the terrace. The upper boundary of the Area is defined as the uppermost edge of this outcrop, and shall be considered a straight line between the visible edges where the outcrop is covered by surface talus. The west end of the Area is defined as where the terrace narrows and merges with a dolerite talus slope on the flank of the north-west ridge of Oliver Peak. The boundary at the west dips steeply from where the upper outcrop disappears, following the border of the dolerite talus with the terrace sandstone down to the western-most corner. The east boundary is defined as the 1 615 m contour, which follows closely the edge of an outcrop which extends much of the width of the terrace. At the southern-most corner of the Area the terrace merges with the slopes into the valley to the east: from this point the boundary extends upward to the 1 700 m contour, from where it follows the line of outcrop defining the south boundary.

3 Map



Note This map predates the current terminology — Site of Special Scientific Interest No. 19 is now Antarctic specially protected area No. 138.

Part 39 Antarctic specially protected area No. 139

1 Name and location


Biscoe Point, Anvers Island.

Latitude 64° 49' S, longitude 63° 49' W.

2 Description

Biscoe Point (latitude 64° 49' S, longitude 63° 49' W) is situated on the southeast side of Biscoe Bay on the south side of Anvers Island in the Palmer Archipelago off the mid-west coast of the Antarctic Peninsula. The site includes the rocky promontory ending in Biscoe Point, the smaller headland immediately to the north and the small islet off the south-west of Biscoe Point. A narrow area of land between the two promontories is included, as is the inshore marine environment within the site boundaries, which are shown on the map.

3 Map



Note This map predates the current terminology — SSSI No. 20 is now ASPA No. 139.

Part 40 Antarctic specially protected area No. 140

1 Name and location

Parts of Deception Island, South Shetland Islands.

62°57' S Latitude, 60°38' W longitude.

2 Description of values to be protected

Deception Island is an active volcano. Recent eruptions occurring in 1967, 1969 and 1970 (Baker *et al.* 1975) altered many of the topographical features of the island and created new, and locally transient, surfaces for the colonisation of plants and other terrestrial biota (Collins 1969, Cameron & Benoit 1970, Lewis Smith 1984*a, b*). There are a number of sites of geothermal activity, some with fumaroles (Smellie *et al.* 2002).

The flora of the island is unique in Antarctic terms, particularly where associated with these geothermal areas, but also because of the recently formed surfaces which provide known-age habitats for the study of colonisation and other dynamic ecological processes by terrestrial organisms (Lewis Smith 1988).

Five small sites around the coast of Port Foster were adopted under Recommendation XIII-8 (ATCM XIII, Brussels, 1985) as Site of Special Scientific Interest No. 21 on the grounds that “Deception Island is exceptional because of its volcanic activity, having had major eruptions in 1967, 1969 and 1970. Parts of the island were completely destroyed, new areas were created, and others were covered by varying depths of ash. Few areas of the interior were unaffected. The island offers unique opportunities to study colonization processes in an Antarctic environment”.

Following a detailed botanical survey of the island in 2002, the values specified in the original designation were reaffirmed and considerably augmented. The survey identified 11 sub-sites of unique botanical interest.

Those interests were:


- The island has the greatest number of rare and extremely rare plant species of any site in the Antarctic. 28 of the 54 mosses recorded on the island, 4 of the 8 liverworts and 14 of the *ca.* 75 lichen are considered to be rare or extremely rare. Appendix A lists the plant species classed as rare or extremely rare in the Antarctic Treaty Area, which occur on Deception Island. These represent 25%, 17% and *ca.* 4% of the total number of mosses, liverworts and lichens, respectively, known from the Antarctic (Aptroot & van der Knaap 1993, Bednarek-Ochyra *et al.* 2000, Ochyra *et al.* in press, Øvstedal & Lewis Smith 2001). 13 species of moss (including two endemics), 2 species of liverwort and 3 species of lichen growing on Deception Island have not been recorded elsewhere in the Antarctic. No other site in the Antarctic is comparable. This suggests that there is a significant deposition of immigrant propagules (by wind and seabirds), particularly of southern South American provenance, over the Antarctic, which becomes established

only where favourable germinating conditions prevail (e.g. the heat and moisture provided around fumaroles) (Lewis Smith 1984*a, b*). Such sites are unique in the Antarctic Treaty Area.

- The more stable geothermal areas, some of which have fumaroles issuing steam and sulphurous gas, have developed bryophyte communities of varying complexity and density, each with a distinct and unique flora. Most of these areas were created during the 1967–70 series of eruptions, but at least one (Mt. Pond) predates that period. Species growing close to active vents are continuously subjected to temperatures between 30°C to 50°C, thereby posing important questions regarding their physiological tolerance.
- Areas of volcanic ash, mudflows, scoria and lapilli deposited between 1967 and 1970 provide unique known-age surfaces. These are currently being colonised by vegetation and other terrestrial biota, allowing the dynamics of immigration and colonisation to be monitored. These areas are unstable and subject to wind and water erosion, so exposing some areas to continual surface change and a cycle of re-colonisation.
- Kroner Lake, the only intertidal lagoon with hot springs in Antarctica, supports a unique community of brackish-water algae.
- Several sites within the Area, unaffected by ash deposits during the 1967–70 eruptions, support long-established mature communities with diverse vegetation and are typical of the older stable ecosystems on the island.
- The largest known stand of Antarctic pearlwort (*Colobanthus quitensis*), one of only two flowering plants in the Antarctic, is located within the Area. After being virtually eradicated by burial in ash during the 1967 eruption, it has recovered and is now spreading at an unprecedented rate within and beyond the original site. This correlates with the current trend in regional climate change, particularly increasing temperature.

3 Map

Figure 1 - Antarctic Specially Protected Area No. 140, Sites of Exceptional Botanical Importance, Deception Island, South Shetland Islands



Part 41 Antarctic specially protected area No. 141

1 Name and location

Yukidori Valley, Langhovde, Lützow-Holmbukta.


Latitude 69° 00' 30" S, longitude 39° 46' 00" E.

2 Description

Yukidori Valley (69° 00' 30" S, 39° 46' 00" E) is situated in the middle part of Langhovde, on the east coast of Lützow-Holm Bay, Continental Antarctica. The Area encompasses 2.0–2.5 km by 1.8 km, located between a tongue of the ice cap and sea at the western end of the Valley. The location of the Area and its boundaries are shown on the attached map. It is described as all the land within the Area bounded by the following coordinates:

- 69° 14' 00" S, 39° 44' 20" E
- 69° 14' 00" S, 39° 48' 00" E
- 69° 15' 00" S, 39° 48' 00" E
- 69° 15' 00" S, 39° 45' 20" E.

3 Map



Part 42 Antarctic specially protected area No. 142

1 Name and location

Svarthamaren, Mühlig-Hofmannfjella, Dronning Maud Land.


Latitude 71° 33' 17" S to 71° 55' 58" S, longitude 5° 09' 12" E to 5° 15' 12" E.

2 Description

The Area is situated in Mühlig-Hoffmannfjella, Dronning Maud Land, stretching from approximately 71° 33' 17" S, 5° 09' 12" E in the north-west to approximately 71° 55' 58" S, 5° 15' 12" E in the south-east. The distance from the ice front is about 200 km. The Area covers approximately 6.4 km², and consists of the ice-free areas of the Svarthamaren nunatak, including the areas in the immediate vicinity of the ice-free areas naturally belonging to the nunatak (i.e. rocks).

The Norwegian field station Tor is located in the Svarthamaren nunatak at latitude 71° 53' S, longitude 5° 10' E. The station, including a 10 m buffer zone around the station buildings, is excluded from the Area.

3 Map



Note This map predates the current terminology — SSSI is now ASPA.

Part 43 Antarctic specially protected area No. 143

1 Name and location

Marine Plain, Mule Peninsula, Vestfold Hills, Princess Elizabeth Land.


Latitude 68° 37' 50" S, longitude 78° 07' 55" E.

2 Description

Marine Plain ASPA lies approximately 10 km south-east of Davis Station in the Vestfold Hills. The Area (23.4 km², 68° 37' 50" S, 78° 07' 55" E) opens into an arm of Crooked Fjord on the southern side of Mule Peninsula, the southern-most of the three major peninsulas that comprise the Vestfold Hills. The Vestfold Hills are a largely ice-free oasis of approximately 512 km² of bedrock, glacial debris, lakes, and ponds, at the eastern side of Prydz Bay, Princess Elizabeth Land.

Starting at the most northerly point of the Area the boundary description is as follows: commencing at 68° 36' 34" S, 78° 09' 28" E, then south-easterly to 68° 36' 45" S, 78° 10' 30" E; then south-easterly to 68° 37' 30" S, 78° 12' 30" E, then south along meridian of longitude 78° 12' 30" E to its intersection with the northern shore of Pineapple Lake; then west along that shore to the edge of the Sørsdal Glacier; then westerly along the northern edge of the Sørsdal Glacier to its intersection with the low water mark of the north eastern shore of Crooked Fjord; then westerly along the low water mark of the northern shore of Crooked Fjord (cutting across the outlet of Burton Lake into Crooked Fjord) to its intersection with the meridian of longitude 78° 03' 00" E; then north along meridian of longitude 78° 03' 00" E to its intersection with the parallel of latitude 68° 37' 30" S, then north-easterly to 68° 36' 56" S, 78° 05' 39" E, then north-easterly to the point of commencement.

3 Map



Part 44 Antarctic specially protected area No. 144

Note The declaration of this area as an Antarctic specially protected area ceases to have effect at the end of 31 December 2010 — see section 5.

1 Name


Chile Bay (Discovery Bay), Greenwich Island, South Shetland Islands.

2 Description

The site comprises two small areas of benthic habitat in Chile Bay located as follows:

- Benthic habitat A — between 50 and 100 m depths and the following coordinates: latitude 62° 28.9' S, longitude 59° 41' 12" W; latitude 62° 29' 3" S, longitude 59° 41' 43" W.
- Benthic habitat B — between 100 and 200 m depths and between the following coordinates: latitude 62° 28.3' S, longitude 59° 40' 15" W; latitude 62° 28.7' S, longitude 59° 40' 47" W.

3 Map



Part 45 Antarctic specially protected area No. 145

1 Name and location

Port Foster, Deception Island, South Shetland Islands

- A: between 62°55' 30" S and 62°56' 12" S, and between 60°37' 00" W and 60°38' 00" W
- B: between 62°57' 12" S and 62°57' 54" S, and between 60°36' 20" W and 60°37' 20" W.

2 Description of values to be protected


Values protected under original designation included the diversity of benthic fauna on two different kinds of sea bottom substrates. The original research about the ecological process of recolonization after volcanic eruption needed protection from the risk of undue interference.

Deception Island is a caldera formed by subsidence of a group of Cenozoic volcanoes superimposed along radial faults. Port Foster is an almost entirely enclosed body of water, receiving large volumes of fresh water during periods of melt. In several places there is geothermal activity.

The area is of exceptional ecological interest because of its actively volcanic character. The two habitat areas are subject to long-term research programs and the purpose in designating them is, as far as possible, to reduce the risk of accidental interference that could jeopardize these scientific investigations.

3 Map

Figure 1. Antarctic Specially Protected Area No. 145, Port Foster, Deception Island, South Shetland Islands.



Part 46 Antarctic specially protected area No. 146

Note The declaration of this area as an Antarctic specially protected area ceases to have effect at the end of 31 December 2010 — see section 5.

1 Name and location


South Bay, Doumer Island, Palmer Archipelago.

Latitude 64° 51' 42" S to 64° 51' 58" S, longitude 63° 34' 00" W to 63° 35' 20" W.

2 Description

Doumer Island lies at the south-west entrance to Neumayer Channel. It is separated from Wiencke Island by the Peltier Channel. South Bay lies on the south coast of Doumer Island. The site consists of a small area of coastal and sub-tidal benthos down to 45 m depth as follows: latitude 64° 51' 42" S to the north, between longitude 63° 34' 00" W and longitude 63° 35' 20" W, and to the south by a diagonal line that starts at a point 100 m north of the Refuge (Sub-base Yelcho) on the southern shore of South Bay and extends to latitude 64° 51' 58" S and longitude 63° 34' 00" W.

3 Map



Note This map predates the current terminology — SSSI No. 28 is now ASPA No. 146.

Part 47 Antarctic specially protected area No. 147

1 Name and location

Ablation Valley-Ganymede Heights, Alexander Island.


Latitude 70° 45' S to 70° 55' S, longitude 68° 21' W to 68° 40' W.

2 Description

Ablation Valley-Ganymede Heights (between latitudes 70° 45' S and 70° 55' S and longitudes 68° 21' W and 68° 40' W, approximately 180 km²) is situated on the east side of Alexander Island, the largest island off the western coast of Palmer Land, Antarctic Peninsula. The Area has a central west-east extent of about 10 km and a north-south extent of about 18 km, flanked to the west by the upper part of Jupiter Glacier, to the east by the permanent ice shelf in George VI Sound, to the north by Grotto Glacier and to the south by the lower reaches of Jupiter Glacier.

The designated Area comprises the entire Ablation Valley-Ganymede Heights massif, bounded in the west by the principal ridge dividing Jupiter Glacier from the main Ablation-Moutonnée-Flatiron valleys. In the east, the boundary is defined by the western margin of George VI Ice Shelf. The northern boundary of the Area is defined as the principal ridge dividing Grotto Glacier from Erratic Valley and other tributary valleys feeding into Ablation Valley, immediately to the south. In the north-west of the Area, the boundary extends across the mostly-glaciated col separating upper Jupiter Glacier from Ablation Valley. The southern boundary of the Area, from east of the principal ridge on the west side of Flatiron Valley to where Jupiter Glacier joins George VI Ice Shelf, is defined as the northern lateral margin of Jupiter Glacier. As the margin between Ablation Lake and George VI Ice Shelf is in places indistinct, the eastern boundary of the Area at Ablation Valley is defined as a straight line extending due south from the eastern extremity of Ablation Point to where the ice shelf abuts land, and from where the eastern boundary follows the land/ice shelf margin. The physiography is similar further south at Moutonnée Lake, and the eastern boundary in this locality is defined as a straight line extending from the eastern extremity of the point on the northern side of (and partially enclosing) Moutonnée Lake to the locality of a prominent meltwater pool where the ice shelf abuts land, and from where the boundary follows the land/ice shelf margin south to where Jupiter Glacier and George VI Ice Shelf adjoin. The Area thus includes the entirety of Ablation and Moutonnée lakes and those parts of the ice shelf behind which they are impounded.

3 Map



Part 48 Antarctic specially protected area No. 148

1 Name and location

Mount Flora, Hope Bay, Antarctic Peninsula.


Latitude 63° 25' S, longitude 57° 01' W.

2 Description

Mount Flora (latitude 63° 25' S, longitude 57° 01' W, 0.3 km²) is situated on the south-eastern flank of Hope Bay, at the northern end of Trinity Peninsula, Antarctic Peninsula. The summit of Mount Flora (520 m) is approximately 1 km from the southern shore of Hope Bay.

The boundaries designated in the original management plan have been revised in the current management plan to include all of the known exposed fossiliferous strata on the northern slopes of Mount Flora. The summit ridge and highest peak of Mount Flora (520 m), which were formerly within the boundary, are comprised of non-fossiliferous volcanic rocks and have now been excluded from the Area. The boundary runs from the north summit of Mount Flora (516 m) — the highest point of the boundary — westward down the ridge to the Kenney Glacier, the eastern margin of Kenney Glacier northward to the 150 m contour, eastward along the 150 m contour to the north-western margin of the Flora Glacier, the north-western margin of the Flora Glacier south-westward to the ridge leading westward to the north summit of Mount Flora. Where present, the glacier margins, lower outcrops, western ridge and northern summit of Mount Flora form visually obvious features that indicate the boundaries: the Area remains otherwise unmarked.

3 Map



Part 49 Antarctic specially protected area No. 149

1 Name and location

Cape Shirreff and San Telmo Island, Livingston Island, South Shetland Islands.

Latitude 62°29' S, longitude 60°48' W.

2 Description of values to be protected


Following designation, the size of the Cape Shirreff Antarctic fur seal colony increased to a level at which biological research could be undertaken without threatening continued increase. A survey of the South Shetland Islands and the Antarctic Peninsula identified Cape Shirreff — San Telmo Island as the most suitable site to monitor Antarctic fur seal colonies that could be affected by fisheries around the South Shetland Islands. In order to accommodate the monitoring program the SPA was redesignated as Site of Special Scientific Interest (SSSI) No. 32 through Recommendation XV-7 (1989) following a joint proposal by Chile, the United Kingdom and the United States of America. Designation was on the grounds that the “presence of both Antarctic fur seal and penguin colonies, and of krill fisheries within the foraging range of these species, make this a critical site for inclusion in the ecosystem monitoring network being established to help meet the objectives of the Convention on the Conservation of Antarctic Marine Living Resources (CCAMLR). The purpose of the designation is to allow planned research and monitoring to proceed, while avoiding or reducing, to the greatest extent possible, other activities which could interfere with or affect the results of the research and monitoring programme or alter the natural features of the Site”. The boundaries were enlarged to include San Telmo Island and associated nearby islets. Following a proposal prepared by Chile and USA, the Area was subsequently designated as CCAMLR Ecosystem Monitoring Program (CEMP) Site No. 2 through CCAMLR Conservation Measure 82/XIII (1994), with boundaries identical to SSSI No. 32.

The current Management Plan reaffirms the exceptional scientific and monitoring values associated with the large and diverse populations of seabirds and pinnipeds which breed within the Area, and in particular those of the Antarctic fur seal colony. The Antarctic fur seal colony is the largest in the Antarctic Peninsula region and is the most southerly that is large enough to study growth, survival, diet, reproduction parameters: it numbered around 20,000 individuals in 2003–2004. Monitoring of the Antarctic fur seal colony began in 1965 (Aguayo and Torres, 1966; 1967) and seasonal data are available from 1991, making this one of the longest continuous Antarctic fur seal monitoring programs. As part of the CEMP, monitoring is established to detect and avoid possible adverse effects of fisheries on dependant species such as pinnipeds and seabirds, as well as target species such as Antarctic krill (*Euphausia superba*). Long-term studies are assessing and monitoring the survival, feeding ecology, growth, condition, reproduction, behavior, vital rates, and abundance of pinnipeds and seabirds that breed within the Area. Data from these studies will be compared alongside environmental and other biological data and fisheries statistics in order to help identify possible cause-effect relationships between fisheries and pinniped and seabird populations.

In 2001–02 imprints of megaflora were discovered in rocks incorporated within moraines of the Livingston Island glacier (Palma-Heldt *et al.*, in review 2004). The fossiliferous rocks are of two different ages, and early interpretations suggest the rocks may assist understanding relationships between nearby fossil localities at Williams Point and Byers Peninsula, also on Livingston Island. Studies of the fossils are on-going and it is important that they are protected from over-sampling.

The original values of the protected area associated with the plant and invertebrate communities cannot be confirmed as primary reasons for special protection of the Area because there is a lack of data available describing the communities.

3 Map



Part 50 Antarctic specially protected area No. 150

Note The declaration of this area as an Antarctic specially protected area ceases to have effect at the end of 31 December 2010 — see section 5.

1 Name and location

Ardley Island, Maxwell Bay, King George Island.


Latitude 62° 13' S, longitude 58° 54' W.

2 Description

Ardley Island (62° 13' S, 58° 54' W) is situated about 500 m east of the coast of Fildes Peninsula, Maxwell Bay, King George Island. It is about 1 km south-east of the Soviet station Bellingshausen and the Chilean station Teniente Marsh, and about 0.5 km east of the Chinese station Great Wall.

The site comprises the entire island and its associated littoral zone, including the isthmus between the island and Fildes Peninsula to the west. The island is about 2.0 km long and 1.5 km at its widest, and rises to about 50 m altitude.

3 Map



Note This map predates the current terminology — SSSI No. 33 is now ASPA No. 150.

Part 51 Antarctic specially protected area No. 151

1 Name

Lions Rump, King George Island, South Shetland Islands.


2 Description

The site is located on the southern coast of King George Bay, King George Island, in the South Shetlands Islands. It is described as all the land and sea falling within the area bounded by the following co-ordinates:

- 62° 07' 48" S, 58° 09' 17" W
- 62° 07' 49" S, 58° 07' 14" W
- 62° 08' 19" S, 58° 07' 19" W
- 62° 08' 16" S, 58° 09' 15" W.

The Area includes the littoral and sublittoral zones extending from the eastern end of Lajkonik Rock to the most northerly point of Twin Pinnacles. From this point the boundary extends to the eastern-most end of the columnar plug of Lions Head to the east of White Eagle Glacier. On land, the Area includes the coast of raised beaches, freshwater pools and streams on the south side of King George Bay, around Lions Cove, and the moraines and slopes which lead to the lower ice tongue of White Eagle Glacier, then westward to a small moraine which protrudes through the ice cap south-east of Sukiennice Hills.

3 Map



Note This map predates the current terminology — SPA is now ASPA.

Part 52 Antarctic specially protected area No. 152

1 Name and location

Western Bransfield Strait off Low Island, South Shetland Islands.

Latitude 63° 15' S to 63° 30' S, longitude 62° 00' W to 62° 45' W.


2 Description

Bransfield Strait is a deep water passage approximately 220 km long and 120 km wide between the Antarctic Peninsula and the numerous islands that comprise the South Shetland Islands. The Drake Passage is to the north and to the west is the Bellingshausen Sea. The Area lies approximately 80 km west of the Antarctic Peninsula, mostly within the 200 m isobath directly south and west of Low Island. Low Island is the southern-most of the South Shetland Islands, lying 60 km south-west of Deception Island and 25 km south-east of Smith Island. To the west and south of Low Island, and for approximately 20 km from the shore, the sea floor slopes gently from the intertidal zone to depths of approximately 200 m. The sea floor slopes steeply to the east of Low Island, reaching depths of up to 1 200 m in this part of Bransfield Strait.

The revised boundaries of the Area at Western Bransfield Strait are defined in the north as the line of latitude at 63° 15' S and in the south at 63° 30' S, in the east the boundary is defined as the line of longitude at 62° 00' W and in the west 62° 45' W. The north-eastern boundary is defined as the shoreline of Low Island, extending from 62° 00' W, 63° 20' S in the south-east (approximately two kilometres from Cape Hooker) to 62° 13' 30" W, 63° 15' S in the north-west (Cape Wallace). The coastline boundary on the western and southern shores of Low Island is defined as the high tide level, and the intertidal zone is included within the Area. The Area extends a maximum of 27.6 km north-south and a maximum of 37.15 km east-west, encompassing an area of approximately 900 km². Boundary markers have not been installed because in the marine area this is impractical, while at Low Island the coast itself is a clearly defined and visually obvious boundary feature.

3 Map

Map 1. ASPA No. 152 Western Bransfield Strait: bathymetric map



Part 53 Antarctic specially protected area No. 153

1 Name and location

Eastern Dallmann Bay off Brabant Island, Palmer Archipelago.

Latitude 63° 53' S to 64° 20' S, longitude 62° 16' W to 62° 45' W.


2 Description

Dallmann Bay (between latitudes 63° 53' S and 64° 20' S and from longitude 63° 16' W eastward to the western shore of Brabant Island) is situated approximately 65 km west of the Antarctic Peninsula, between Brabant Island and Anvers Island, with Bransfield Strait to the north and Gerlache Strait to the south.

The designated Area is defined in the south by latitude 64° 20' S, extending from Fleming Point westward for two kilometres to 62° 40' W. From this location, the western boundary extends due north on longitude 62° 40' W for 18.5 km to 64° 10' S, south-south-west of Astrolabe Needle. The western boundary then extends north-north-west almost 19 km to 62° 45' W, 64° 00' S. The western boundary then extends approximately 13 km due north on longitude 62° 45' W to latitude 63° 53' S, the northern boundary of the Area. The northern boundary extends along latitude 63° 53' S from 62° 45' W to 62° 16' W, being a distance of approximately 23.4 km. The eastern boundary extends due south approximately 16 km from 62° 16' W, 63° 53' S to the eastern extremity of Pasteur Peninsula, Brabant Island, at 62° 16' W, 64° 02' S. From there, the eastern boundary is defined as the mean high water mark of the northern and western coastline of Brabant Island, which includes the intertidal zone within the Area. The Area is 50 km from north to south and extends up to a maximum of 23.4 km east-west. West of Brabant Island the width of the Area ranges between 10 km (at Guyou Bay) and 1.5 km (near Claude Point). The total area is approximately 580 km².

3 Map

Map 1. ASPA No. 153 Eastern Dallmann Bay: bathymetric map



Part 54 Antarctic specially protected area No. 154

1 Name and location

Botany Bay, Cape Geology, Victoria Land.


Latitude 77° 00' 14" S, longitude 162° 32' 52" E.

2 Description

Cape Geology is situated in the south-western corner of Granite Harbour, southern Victoria Land, at 77° 00' 14" S, 162° 32' 52" E approximately 100 km north-west of Ross Island.

The north-west corner of the Area is marked by a brass plaque in a boulder (M1, 2 m) 400 m south-west of Cape Geology. The west boundary is defined by a line extending first 260 m south-south-east from M1 to a large boulder (marked by a cairn) with terrier bolt (M2) at an elevation of 118 m on the ridge above the campsite; thence the boundary extends 250 m up this ridge to a point at 162 m elevation marked by an iron tube with bamboo pole. The west boundary extends a further 300 m up this ridge to a large pointed rock at 255 m elevation near the edge of the permanent ice field. The boundary then extends 150 m south across the ice field to the west edge of a prominent line of exposed rock and moraine in the south-west corner of the Area at 325 m elevation. The south boundary follows this line of rock east until the exposure is buried by the ice field, thence south-east across the ice field for 500 m to the edge of a second and more prominent exposure at an elevation of just over 400 m (M3). The boundary follows the upper edge of this exposure and then crosses the ice field south-east to an elevation of approximately 325 m where the ice-free eastern boundary ridge and the ice field converge. The east boundary follows the ridge crest for 1 550 m in a north-east direction to a large pointed rock on the ridge (M4, 392 m) where the east boundary turns to descend due north to the coast at the eastern extremity of the boulder beach of Botany Bay (M5, 5 m). The mean high water mark of the coastline of Botany Bay and Cape Geology forms the northern boundary of the Area.

3 Map



Part 55 Antarctic specially protected area No. 155

1 Name and location

Cape Evans, Ross Island.

77°38' S Latitude, 166°24' E longitude.

2 Description of values to be protected


The significant historic value of this Area was formally recognised when it was listed as Historic Site and Monument Nos 16 and 17 in Recommendation 9 (1972). An area containing both sites was designated as Specially Protected Area No. 25 in Measure 2 (1997) and redesignated as Antarctic Specially Protected Area 155 in Decision 1 (2002).

The *Terra Nova* hut (Historic Site and Monument No. 16) is the largest of the historic huts in the Ross Sea region. It was built in January 1911 by the British Antarctic *Terra Nova* Expedition of 1910–1913, led by Captain Robert Falcon Scott, RN. It was subsequently used as a base by the Ross Sea party of Sir Ernest Shackleton's Imperial Trans-Antarctic Expedition of 1914–1917.

Some of the earliest advances in the study of earth sciences, meteorology, flora and fauna are associated with the *Terra Nova* Expedition based at this site. The data collected can provide a bench mark against which to compare current measurements. The history of these activities and the contribution they have made to the understanding and awareness of Antarctica therefore contribute to both the historic and scientific value of the site.

The Cape Evans site is one of the principal sites of early human activity in Antarctica. It is an important symbol of the Heroic Age of Antarctic exploration, and as such, has considerable historical significance.

3 Map



Part 56 Antarctic specially protected area No. 156

1 Name and location


Lewis Bay, Mount Erebus, Ross Island.

Latitude 77° 25' 29" S, longitude 167° 28' 30" E.

2 Description

The designated Area on Ross Island encompasses the crash zone (centred on 77° 25' 29" S, 167° 28' 30" E, elevation 520 m (1 720 feet)) and the surrounding glacial ice 2 km above and to either side of this position, extends as a 4 km wide 'rectangle' down to the sea, and includes the airspace above this region to an altitude of 1 000 m (3 280 feet) with the exception of a 200 m wide air access 'corridor' along the coastline. The west boundary of the Area is the 167° 23' 33" E meridian; the east boundary is the 167° 33' 27" E meridian. The south boundary is the 77° 26' 33" S parallel, while the north boundary is defined by the coastline.

3 Map



Part 57 Antarctic specially protected area No. 157

1 Name and location

Backdoor Bay, Cape Royds, Ross Island.

77°33' 10.7" S Latitude, 166°10' 6.5" E longitude.

2 Description of values to be protected


The significant historic value of this Area was formally recognised when it was listed as Historic Site and Monument No. 15 in Recommendation 9 (1972). It was designated as Specially Protected Area No. 27 in Measure 1 (1998) and redesignated as Antarctic Specially Protected Area (ASPA) 157 in Decision 1 (2002).

The hut on which this Area is centred was built in February 1908 by the British Antarctic (*Nimrod*) Expedition of 1907–1909 which was led by Sir Ernest Shackleton. It was also periodically used by the Ross Sea Party of Shackleton's Imperial Trans-Antarctic Expedition of 1914–1917.

Structures associated with the hut include stables, kennels, a latrine and a garage created for the first motor vehicle in Antarctica. Other significant relics in the Area include an instrument shelter, supply depots, and a rubbish site. Numerous additional artefacts are distributed around the Area.

Cape Royds is one of the principal areas of early human activity in Antarctica. It is an important symbol of the Heroic Age of Antarctic exploration and, as such, has considerable historical significance. Some of the earliest advances in the study of earth sciences, meteorology, flora and fauna in Antarctica are associated with the *Nimrod* Expedition which was based at this site. The history of these activities and the contribution they have made to the understanding and awareness of Antarctica give this Area significant scientific, aesthetic and historic values.

3 Map



Part 58 Antarctic specially protected area No. 158

1 Name and location

Hut Point, Ross Island.

Latitude 77° 50' 50" S, longitude 166° 38' E.


2 Description of values to be protected

The significant historic value of this Area was formally recognised when it was designated as Historic Site and Monument No. 18 in Recommendation 9 (1972). It was designated as Specially Protected Area No. 28 in Measure 1 (1998) and redesignated as Antarctic Specially Protected Area 158 in Decision 1 (2002).

The hut was built in February 1902 during the National Antarctic (*Discovery*) Expedition of 1901–1904, led by Captain Robert Falcon Scott who later found it a valuable advance staging point for journeys on the “Barrier” during his 1910-1913 expedition. It was also used by Sir Ernest Shackleton during the 1907-1909 British Antarctic Expedition and later by his stranded Ross Sea Party during the Imperial Trans-Antarctic Expedition of 1914–1917. This building was prefabricated in Australia to an ‘outback’ design with verandahs on three sides.

The Hut Point site is one of the principal sites of early human activity in Antarctica. It is an important symbol of the Heroic Age of Antarctic exploration and, as such, has considerable historical significance. Some of the earliest advances in the study of earth sciences, meteorology, flora and fauna in Antarctica are associated with the *Discovery* Expedition based at this site. The history of these activities and the contribution they have made to the understanding and awareness of Antarctica give this Area significant scientific, aesthetic and historic values.

3 Map



Part 59 Antarctic specially protected area No. 159

1 Name and location

Cape Adare, Borchgrevink Coast, Northern Victoria Land.

Eastern longitudes 11° 22' 40" and 11° 54' 20".

Southern latitudes 70° 43' 50" and 70° 46' 40".

2 Description of values to be protected

The historic value of this Area was formally recognized when it was listed as Historic Site and Monument No. 22 in Recommendation VII-9 (1972). It was designated as Specially Protected Area No. 29 in Measure 1 (1998) and redesignated as Antarctic Specially Protected Area (ASPA) 159 in Decision 1 (2002).


There are three main structures in the Area. Two were built in February 1899 during the British Antarctic (*Southern Cross*) Expedition led by C. E. Borchgrevink (1898–1900). One hut served as a living hut and the other as a store. They were used for the first winter spent on the Antarctic continent.

Scott's Northern Party hut is situated 30 metres to the north of Borchgrevink's hut. It consists of the collapsing remains of a third hut built in February 1911 for the Northern Party led by V. L. A. Campbell of R. F. Scott's British Antarctic (*Terra Nova*) Expedition (1910–1913), which wintered there in 1911.

In addition to these features there are numerous other historic relics located in the Area. These include stores depots, a latrine structure, two anchors from the ship *Southern Cross*, an ice anchor from the ship *Terra Nova*, and supplies of coal briquettes. Other historic items within the Area are buried in guano.

Cape Adare is one of the principal sites of early human activity in Antarctica. It is an important symbol of the Heroic Age of Antarctic exploration and, as such, has considerable historical significance. Some of the earliest advances in the study of earth sciences, meteorology, flora and fauna in Antarctica are associated with the two earliest expeditions based at this site. The history of these activities and the contribution they have made to the understanding and awareness of Antarctica give this Area significant scientific, aesthetic and historic values.

3 Map



Part 60 Antarctic specially protected area No. 160

1 Name and location

Frazier Islands, Windmill Islands, Wilkes Land, East Antarctica.


Latitude 66° 14' S, longitude 110° 10' E.

2 Description

The Frazier Islands are located at latitude 66° 14' S, longitude 110° 10' E. The three islands (Nelly, Dewart and Charlton Island) lie in the eastern part of Vincennes Bay approximately 16 km to the west-north-west of Casey Station.

The ASPA comprises the entire terrestrial area of the three islands, with the seaward boundary at the low water mark. The total area of the Frazier Islands ASPA is approximately 0.6 km². There are no boundary markers.

3 Map



Part 61 Antarctic specially protected area No. 161

1 Name and location

Terra Nova Bay, Ross Sea.

Latitude 74° 42' 57" S to 74° 48' 00" S, longitude 164° 10' 00" E.


2 Description

The designated Area is situated in Terra Nova Bay, between the Campbell Glacier Tongue and Drygalski Ice Tongue, Victoria Land. The Area is confined to a narrow strip of coastal waters to the south of Terra Nova Bay Station (Italy), extending approximately 9.4 km in length and generally within 1.5–7 km of the shore, comprising an area of 29.4 km².

The western boundary of the Area is defined as the mean high water mark along the coastline extending between 74° 42' 57" S in the north (2.3 km south of Terra Nova Bay Station) and 74° 48' 00" S in the south (the southern shore of Adélie Cove), and includes the intertidal zone. The northern boundary of the Area is defined as the 74° 42' 57" S line of latitude, extending from the coast 1.55 km eastward to the 164° 10' 00" E line of longitude. The boundary position may be recognised near the shore by the presence of a large and distinctive offshore rock in the northernmost cove on the coast south of Terra Nova Bay Station, which is a unique feature on this stretch of coast. The southern boundary is defined as the 74° 48' 00" S line of latitude, extending from the coast 3.63 km eastward to

the 164° 10' 00" E line of longitude. The boundary position may be recognised visually as being at the southern shore of the mouth of Adélie Cove, immediately south of a distinctive rocky outcrop at the base of the coastal cliffs. The eastern boundary of the Area is defined as the 164° 10' 00" E line of longitude extending between 74° 42' 57" S in the north and 74° 48' 00" S in the south.

3 Map



Note Part 62 is reserved.

Part 63 Antarctic Specially Protected Area No. 163

1 Name and Location

Dakshin Gangotri Glacier, Dronning Maud Land.

Latitude 70°43' 50" S, longitude 70°46' 40" E.

2 Description of values to be protected

Historic value

Dakshin Gangotri Glacier is a small tongue of polar continental ice sheet, overriding the Schirmacher Oasis of central Dronning Maud Land (CDML). It was identified by the second Indian Antarctic Expedition in 1983 and since then its snout is being monitored continuously.

Scientific value


With the availability of this vast amount of data for the past two decades, it has become a valuable site for observing the changes in the movement of the Antarctic ice sheet under the impact of global warming. The area has primary scientific importance for glaciologists and environmental scientists. Due to the scientific values of the Area and the nature of the research, it is protected as an Antarctic Specially Protected Area consistent with Articles 2, 3, 5 and 6 of Annex V of the Protocol on Environmental Protection to the Antarctic Treaty, to prevent interference with ongoing planned scientific investigations.

Environmental value

In the designated Area exploration shows the faunal diversity and ecology of the moss-inhabiting terrestrial invertebrate fauna and it is also extensively explored lichenologically. Schirmacher Oasis is also important area for the Algal and cyanobacterial flora diversity. Terrestrial mosses are quite widespread in the Schirmacher Oasis colonizing a range of habitats. The bryophytes, because of their poikilohydric nature and alternative strategy of adaptation, are one of the very few plant groups which grow in Antarctica. As such their role in habitat modification, nutrient cycling, primary production and providing shelter and security to associated invertebrate animals, for example the bryobionts, bryophiles, bryoxenes assume a particular significance. Bryophytes have been reported in Schirmacher Oasis (divided in Eastern, Central and Western Schirmacher). Lichens, fungi, algae and bacteria have also been reported in the Area. Distribution of algae and cyanobacteria flora of fresh water streams of the Oasis at the designated area are studied. Examination of algae and cyanobacteria is conducted in the glacier-melt water stream. The species reported are *G. magma*, *Chaemosiphon subglobosus*, *Oscillatoria limosa*, *O. limnetica*, *P. frigidum*, *P. autumnale*, *Nostoc commune*, *N. punctiforme*,

Calothrix gracilis, *C. brevissima*, *Uronema* sp., and *Cosmarium leave*. Among the cyanobacteria encountered in the stream of Schirmacher Oasis, contribution by N₂-fixing species might play a significant role in nitrogen economy of the ecosystem through N₂-fixation. Studies on polar Skuas are also conducted at Schirmacher Oasis and their nesting and breeding success is reported around the designated place.

3 Map



Part 64 Antarctic Specially Protected Area No. 164

1 Name and Location

Scullin and Murray Monoliths, MacRobertson Land, East Antarctica.

Latitude 67° 47' 01" S, longitude 66° 40' 31" E.

2 Description of values to be protected

Scullin and Murray Monoliths (67° 47' S, 66° 42' E and 67° 47' S, 66° 53' E) hold the greatest concentration of breeding seabird colonies in East Antarctica, including the second largest colony of Antarctic petrels *Thalassoica antarctica*. The Scullin and Murray Monoliths ASPA is a breeding locality for at least 160 000 pairs of Antarctic petrels from a minimum estimated global total of approximately half a million pairs (van Franeker *et al.* 1999). Scullin Monolith was recently recognised as a candidate Important Bird Area (IBA) using IUCN/BirdLife International criteria on known breeding populations by the SCAR Bird Biology subcommittee in mid-2002 (SCAR unpubl. data).

Adélie penguin colonies occupy the lower slopes of both monoliths, extending almost to the foreshore. Approximately 50 000 pairs nest on Scullin Monolith and a further 20 000 pairs on Murray Monolith. This

represents approximately 10% of the Adélie penguin breeding population for East Antarctica and approximately 3% of the global population.

Many of the ocean-facing slopes of both monoliths are used for breeding by petrels. Extensive breeding colonies of four species of petrels occupy many of the steeper, higher-altitude slopes of both monoliths. The Antarctic petrel colony on Scullin Monolith is second in population only to the colony at Svarthameren in the Mühlig Hofmannfjella, in Dronning Maud Land. South Polar skuas nest throughout the ASPA, making use of the high density of breeding seabirds as prey during their breeding season.

While larger colonies of seabirds are known from elsewhere in East Antarctica (e.g. the Rauer Group), it is the extensive populations (total known breeding population conservatively estimated at 230 000 pairs equivalent to a minimum of 460 000 individual breeding seabirds) and rich species diversity (seven breeding species) within the very small ice-free areas of Scullin and Murray Monoliths (estimated ice free area of 1.9 and 0.9 km², respectively, total of 2.8 km²) that make these the greatest concentration and one of the most diverse seabird breeding localities known from East Antarctica.

There are no data on population trends available, and the census and survey data collected in 1986/87 serve as baseline data for all future ornithological work in the Area. Some limited census data were collected from Reference Breeding Groups (RBGs) established in the mid 1980s to monitor the Antarctic petrel population; there have been no surveys of these RBGs for more than a decade. Many Adélie penguin breeding populations throughout East Antarctica have increased in the last 20 or so years, and it is likely that the Adélie penguin population within the Scullin and Murray Monoliths ASPA is greater than the 70 000 pairs reported from 1986/87. Further, it is likely that the 1986/87 census under-estimated the breeding population of Antarctic petrels, given the census occurred late in the breeding season.

Aesthetic and wilderness values

In addition to the outstanding ecological and scientific values already identified, the Area possesses outstanding aesthetic values in the geomorphology of the two Monoliths and the spectacular nature of the glaciers descending from the Continental plateau that flow around the Monoliths ending in calving glaciers. The near-vertical aspects of both Monoliths dropping to the sea, used by a high number of seabirds for nesting, represent an unique landscape in the Antarctic. The very large breeding assemblage of undisturbed seabirds in a setting of high aesthetic and wilderness values warrants the highest level of protection.

Human history


Recorded visits to Scullin and Murray Monoliths are few. Scullin and Murray Monoliths were first visited during the second BANZARE voyage in 1930-31, on 13th February 1931 (Grenfell Price 1962, Fletcher 1984). A brief landing was made at Scullin Monolith on 26 February 1936 from the R.R.S. *William Scoresby*, when an ascent was made to a height of several

hundred metres (Rayner 1940). A landing by the Norwegian, Lars Christensen was made on 30 January 1937, when Scullin Monolith was visited (Christensen 1938, 1939). Australian National Antarctic Research Expeditions (ANARE) personnel have made few visits to the Area from Mawson station, approximately 160 km to the west. The only recorded stay within the Area was a six-day visit (1–6 February 1987), when comprehensive ornithological surveys were conducted (Alonso *et al.* 1987). A fibreglass ‘Apple’ refuge was established within the ASPA for this visit, and as of 13 October 2002, was intact. The first visit by a commercial tourist vessel to the Area was made on 10 December 1992, when passengers were landed at Scullin and Murray Monoliths. Brief tourist visits were made to Scullin Monolith on 7 December 1997, to Scullin and Murray Monoliths on 8 January 1998 and, Scullin and Murray Monoliths on 18 December 2002. Compared to many sites in East Antarctica, Scullin and Murray Monoliths have been visited infrequently, and with the one known exception, all visits have been brief (less than a day). Further, with little activity conducted during those visits, the Area, and in particular the avifauna, is of particular value as a relatively undisturbed area suitable for use in the future as a reference site for other areas that experience a greater level of human visitation and extent of activities.

Nomenclature

Mawson named both Monoliths during the second BANZARE voyage. Murray Monolith was named after Sir George Murray, Chief Justice of South Australia, Chancellor of the University of Adelaide and a patron of the Expedition, while Scullin Monolith was named after James H. Scullin, Prime Minister of Australia from 1929–31.

3 Map



Schedule 2 Antarctic specially managed areas

Note Part 1 is reserved.

Part 2 Antarctic Specially Managed Area No. 2

1 Name and location

McMurdo Dry Valleys, southern Victoria Land.

The Area boundaries have been defined primarily on the basis of the hydrological catchments in the McMurdo Dry Valleys, including all of the ice-free ground and adjacent areas within these catchments, all of the Convoy Range, and the catchment of the Alph River. Starting at the northwest corner and moving counter-clockwise, the boundary is delineated by the following features: the northwest tip of Allan Nunatak (76.7167° S, 159.6667° E), Carapace Nunatak (76.8833° S, 159.4° E), Mt. DeWitt (77.2° S, 159.8333° E), the western edge of Horseshoe Mountain (77.5667° S, 159.95° E), Depot Nunatak (77.75° S, 160.0667° E), the southernmost peak of the Lashly Mountains (77.9606° S, 159.5603° E), Mt. Kempe (78.3167° S, 162.7167° E), the Pyramid (78.35° S, 163.5° E), the east side of Heald Island (78.25° S, 163.8167° E), De Master Point (off the eastern end of Marshall Valley, 78.0792° S 164.4131° E), north along the coast following the mean low tide level to the east side of Tripp Island (76.6333° S, 162.7° E), the southern edge of Fry Glacier (76.6333° S, 162.3° E), and, again the northwest tip of the Allan Nunatak (76.7167° S, 159.6667° E).

2 Description of values to be protected

The McMurdo Dry Valleys are characterized as the largest relatively ice-free region in Antarctica with approximately thirty percent of the ground surface largely free of snow and ice. The region encompasses a cold desert ecosystem, whose climate is not only cold and extremely arid (in the Wright Valley the mean annual temperature is -19.8°C and annual precipitation is less than 100 mm water equivalent), but also windy. The landscape of the Area contains glaciers, mountain ranges, ice-covered lakes, meltwater streams, arid patterned soils and permafrost, sand dunes, and interconnected watershed systems. These watersheds have a regional influence on the McMurdo Sound marine ecosystem. The Area's location, where large-scale seasonal shifts in the water phase occur, is of great importance to the study of climate change. Through shifts in the ice-water balance over time, resulting in contraction and expansion of hydrological features and the accumulations of trace gases in ancient snow, the McMurdo Dry Valley terrain also contains records of past climate change. The extreme climate of the region serves as an important analogue for the conditions of ancient Earth and contemporary Mars, where such climate may have dominated the evolution of landscape and biota.


The Area is characterized by unique ecosystems of low biodiversity and reduced food web complexity. However, as the largest ice-free region in Antarctica, the McMurdo Dry Valleys also contain relatively diverse habitats compared with other ice-free areas. The Area contains unusual microhabitats and biological communities (such as endolithic and Cryoconite systems) as well as special geological features and minerals (for example, salt deposits and desert pavements). Some of these special geological features are of value because they contain an extremely long record of natural events. The long-term data sets for environmental observations that have been collected in this region are some of the longest in Antarctica. The McMurdo Dry Valleys contain indicators of past and present regional climate change, as well as features that play a role in influencing local climate change.

These scientific values are also of global as well as regional importance. The Area is a valuable resource for understanding landscape processes and the stability of Antarctic ice sheets. The McMurdo Dry Valleys contain unique surface deposits including glacially deposited and modified sediments, sand dunes, desert pavement, glacio-lacustrine sediments, and marine fjord sediments containing valuable records of planetary change. The soil, rock, water, and ice environments and their associated biota are of scientific value as model ecosystems that allow deep insights into natural processes operating throughout the biosphere. Finally, the species that reside in the McMurdo Dry Valleys provide a biological resource for understanding adaptation to extreme environments, and are true end members of ecological continua.

The McMurdo Dry Valleys are also valued for their wilderness quality. They represent a nearly pristine environment largely undisturbed and uncontaminated by humans. The dramatic landscape, composed of high ridges and sweeping valleys, and contrasts of ice-free and glacier-covered terrain creates unique vistas with high aesthetic value.

Activities conducted in the area include a variety of scientific research, operations in support of science, media, arts, education and other official national programme visitors; and tourism. A Long Term Ecological Research site has been established in the Taylor Valley.

Map A: Map of the McMurdo Dry Valleys Area



Part 3 Antarctic Specially Managed Area No. 3

1 Name and location

Cape Denison, Commonwealth Bay, George V Land.

Latitude 67° 00' 13" S–67° 00' 50" S, longitude 142° 40' 00.1" E–
142° 41' 27" E.

2 Description of values to be protected

Primary values

This ASMA is proposed on the grounds that Cape Denison is a site of historic, archaeological, social and aesthetic values.

Historic value

Antarctica's 'heroic age' was a period of great human adventure and discovery. Cape Denison, Commonwealth Bay provides the setting for the buildings, structures and relics of the Main Base of the Australasian Antarctic Expedition (AAE) of 1911–14, led by Dr Douglas Mawson.

The prime focus of Mawson's was scientific research. Nevertheless, the expedition also had an exploratory agenda, with the aim of charting the entire Antarctic coastline immediately south of Australia. For this purpose at least five sledging expeditions were undertaken from Cape Denison from spring 1912, including the infamous Far-Eastern Sledging Party during which expeditioners Belgrave Ninnis and Xavier Mertz perished and Mawson himself barely survived. Overall, more than 6 500 km of coastline and hinterland was explored by sledging parties of the Expedition.

Cape Denison contains numerous relics relating to the work of Mawson's expedition, including Mawson's Huts and other significant and relatively untouched artefacts from the 'heroic age'. While the majority is concentrated in the westernmost valley and its immediate surrounds, the historical boundaries of the Main Base extend further. Artefacts and other evidence of occupation, such as food caches, extend across the entire Cape, forming a rich resource of material available for research and interpretation, and potentially yielding scientific data and information about aspects of expeditioner life not included in official written accounts.

Aesthetic values

This ASMA is designated to preserve not only the artefacts remaining *in situ* but also the cultural landscape of Cape Denison in which Mawson and his men lived and worked. Cape Denison is characterised by its almost incessant blizzard conditions, which severely limit access to the region and activities at the site. System and katabatic winds pour down the plateau and funnel through the Cape's valleys; blasting the hut with gusts that in May 1912 reached 322 km/h. (The average wind speed for the month was 98 km/h). Cape Denison is not only the windiest place in Antarctica, but also

the windiest place on Earth at sea level. The site thus provides the physical and symbolic context of the extreme isolation and harsh conditions endured by the expedition members and, by association, all other 'heroic age' researchers and explorers. In designating the entire area as an ASMA, Cape Denison's unique 'sense of place' is protected, with Mawson's Huts and Boat Harbour as the focus of the visual catchment. Mawson's Huts themselves are provided with additional protection in ASPA 162.

Educational values

Cape Denison's wildlife and undisturbed artefacts, framed against the dramatic backdrop of the Antarctic Plateau, represent significant educational values. The Area's isolation and extreme weather provide visitors with a unique insight into the conditions endured by 'heroic age' researchers and explorers, and a chance to form a deeper appreciation of their achievements.

Environmental values

The paucity of relatively ice-free areas in the immediate region means that Cape Denison represents an important assemblage of life forms. The closest ice-free areas of equal or greater size to Cape Denison are approximately 20 km to the east of Cape Denison (from the centroid of the ASMA), and approximately 60 km to the west respectively. A haul-out site for Weddell, leopard and elephant seals, the Cape is also an important breeding area for Adélie penguins, Wilson's storm-petrels, snow petrels and south polar skuas.

Flora at Cape Denison is represented by 13 lichen species distributed on boulders and other moraines throughout the peninsula. No bryophytes are evident. The lichens' distribution on rocks, which are subject to different patterns of snow ablation, makes them vulnerable to trampling and other interference by visitors, however infrequent visitation may be.

Cape Denison has 13 small lakes. These are associated with glacial action, are a permanent feature, and are frozen over for most of the year. Since such lakes are also susceptible to physical, chemical and biological modification within their catchment boundaries, a catchment-based approach to the management of human activities is required.


Scientific values

Mawson, a geologist, planned his expedition in order to examine the theories about continental connection and the processes of glaciation and climate. He also sought to study the South Magnetic Pole and magnetic charting for navigational purposes; to conduct biological studies, including the identification of new species; and to establish a weather station.

Cape Denison provides opportunities to repeat Mawson's experiments and conduct further research into magnetism, meteorology, biology, and other sciences. For example, although Antarctic lakes are generally recognised as valuable due to their relatively simple natural ecosystems, the lakes at Cape Denison have neither been sampled nor their biota studied. There are also numerous non-marine algae present; however, no surveys have been undertaken.

The records from Mawson's expedition provide a dataset against which the results of modern research may be compared, and the site's isolation lends it considerable value for future use as a reference site for other areas that experience a greater level of human activities.

3 Map



Part 4 Antarctic Specially Managed Area No. 4

1 Name and location

Deception Island, South Shetland Islands.

Latitude 62° 57' S, longitude 60° 38' W.

2 Description of values to be protected

i. Natural value

- Deception Island is one of only two volcanoes in the Antarctic at which eruptions have been observed. It was responsible for numerous ash layers dispersed across the South Shetland Islands, Bransfield Strait and the Scotia Sea. Ash from the island has even been recorded in an ice core at the South Pole. The volcano erupted during two short periods during the 20th century, most recently between 1967–1970. It contains a restless caldera that is actively deforming. It is therefore likely that Deception Island will witness further eruptions in the future.
- The Area has an exceptionally important flora, including at least 18 species which have not been recorded elsewhere in the Antarctic. No other Antarctic area is comparable. Of particular importance are the very small, unique biological communities associated with the island's geothermal areas, and the most extensive known community of the flowering plant Antarctic pearlwort (*Colobanthus quitensis*).
- Eight species of seabird breed on the island, including the world's largest colony of chinstrap penguins (*Pygoscelis antarctica*).
- The benthic habitat of Port Foster is of ecological interest due to the natural perturbations caused by volcanic activity.

ii. Scientific value and activities

- The Area is of outstanding scientific interest, in particular for studies in geoscience and biological science. It offers the rare opportunity to study the effects of environmental change on an ecosystem, and the dynamics of the ecosystem as it recovers from natural disturbance.
- Long term seismological and biological data-sets have been collected at Decepción Station (Argentina) and Gabriel de Castilla Station (Spain).

iii. Historic value

- The Area has had a long history of human activity since c.1820, including exploration, sealing, whaling, aviation and scientific research, and as such has played a significant role in Antarctic affairs.
- At Whalers Bay, the Norwegian Hektor whaling station, the cemetery and other artefacts, some of which pre-date the whaling station, are the most significant whaling remains in the Antarctic. The British 'Base B', which was established in the abandoned whaling station, was the first base of the secret World War II expedition 'Operation Tabarin', the forerunner to the British Antarctic Survey. As such, it was one of the earliest permanent research stations in Antarctica. The whalers remain and Base B are listed as Historic Site and Monument (HSM) No. 71.

- The remains of the Chilean Presidente Pedro Aguirre Cerda Station at Pendulum Cove are listed as HSM No. 76. Meteorological and volcanological studies were undertaken at the base from 1955 until its destruction by volcanic eruptions in 1967 and 1969.

iv. Aesthetic value

- Deception Island's flooded caldera, its 'horse-shoe' shape and linear glaciated eastern coastline, its barren volcanic slopes, steaming beaches and ash-layered glaciers provide an unique Antarctic landscape.

v. Educational and Tourism activities

- Deception Island is the only place in the world where vessels can sail directly into the centre of a restless volcanic caldera, providing the opportunity for visitors to learn about volcanoes and other aspects of the natural world, as well as early Antarctic exploration, whaling and science. Deception Island is also one of the most frequently visited sites in Antarctica by tourists.

3

Map




Figure 1. The location of Deception Island in relation to the South Shetland Islands and the Antarctic Peninsula