

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Abamectin	
Sum of avermectin B1a, avermectin B1b and (Z)-8,9 avermectin B1a, and (Z)-8,9 avermectin B1b	
Adzuki bean (dry)	T*0.002
Almonds	T*0.01
Apple	0.01
Avocado	T0.05
Blackberries	T0.1
Blueberries	T*0.02
Cattle, edible offal of	0.1
Cattle fat	0.1
Cattle meat	0.005
Cattle milk	0.02
Chervil	T0.5
Citrus fruits	0.02
Common bean (dry) (navy bean)	T*0.002
Coriander (leaves, stem, roots)	T0.5
Cotton seed	*0.01
Cucumber	0.02
Currant, black	0.02
Egg plant	0.02
Goat fat	0.1
Goat kidney	0.01
Goat liver	0.05
Goat milk	0.005
Goat muscle	0.01
Grapes	0.02
Herbs	T0.5
Hops, dry	0.1
Kaffir lime leaves	T0.5
Lemon grass	T0.5
Lettuce, head	0.05
Lettuce, leaf	T1
Maize	T*0.01
Melons, except watermelon	T0.02
Mung bean (dry)	T*0.002
Mushrooms	T0.05
Onion, Welsh	T0.05
Papaya (pawpaw)	T0.1
Passionfruit	T0.2
Peanut	T*0.002
Pear	0.01
Peas	T0.5
Peppers	T0.1
Pig kidney	0.01
Pig liver	0.02
Pig meat (in the fat)	0.02
Popcorn	T*0.01
Potato	T0.01
Raspberries, red, black	T0.1
Rhubarb	T0.05
Shallot	T0.05
Sheep, edible offal of	0.05
Sheep meat (in the fat)	0.05
Soya bean (dry)	*0.002
Spring onion	T0.05
Squash, Summer	0.02
Strawberry	0.1
Sweet corn (corn-on-the-cob)	T0.05
Tomato	0.05
Watercress	T0.5
Watermelon	T0.02

Acephate	
Acephate (Note: the metabolite methamidophos has separate MRLs)	
Banana	1
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	5
Citrus fruits	5
Cotton seed	2
Edible offal (mammalian)	0.2
Eggs	0.2
Lettuce, head	10
Lettuce, leaf	10
Macadamia nuts	*0.1
Meat (mammalian) [except sheep meat]	0.2
Peppers, Sweet	5
Potato	0.5
Sheep meat	*0.01
Soya bean (dry)	1
Sugar beet	0.1
Tomato	5
Tree tomato (tamarillo)	0.5
Acequinocyl	
Sum of acequinocyl and its metabolite 2-dodecyl-3-hydroxy-1,4-naphthoquinone, expressed as acequinocyl	
Citrus fruits	0.2
Grapes	1.6
Acetamiprid	
Commodities of plant origin: Acetamiprid Commodities of animal origin: Sum of acetamiprid and N-demethyl acetamiprid ((E)-N ¹ -[6-chloro-3-pyridyl)methyl]-N ² -cyanoacetamidine), expressed as acetamiprid	
Citrus fruits	0.5
Cotton seed	*0.05
Cranberry	0.6
Cucumber	T0.2
Date	T5
Edible offal (mammalian)	*0.05
Eggs	*0.01
Grapes	0.35
Meat (mammalian)	*0.01
Milks	*0.01
Potato	*0.05
Poultry, edible offal of	*0.05
Poultry meat	*0.01
Stone fruits [except plums]	1
Tomato	T0.1
Acibenzolar-S-methyl	
Acibenzolar-S-methyl and all metabolites containing the benzo[1,2,3]thiadiazole-7-carboxyl moiety hydrolysed to benzo[1,2,3]thiadiazole-7-carboxylic acid, expressed as acibenzolar-S-methyl	
Cotton seed	*0.02
Edible offal (mammalian)	*0.02
Eggs	*0.02
Meat (mammalian)	*0.02
Milks	*0.005
Poultry, edible offal of	*0.02
Poultry meat	*0.02

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Acifluorfen Acifluorfen	
Edible offal (mammalian)	0.1
Eggs	*0.01
Legume vegetables	0.1
Meat (mammalian)	*0.01
Milks	*0.01
Peanut	0.05
Poultry, edible offal of	0.1
Poultry meat	*0.01
Pulses	0.1
Albendazole Sum of albendazole, its sulfoxide, sulfone and sulfone amine, expressed as albendazole	
Cattle, edible offal of	*0.1
Cattle meat	*0.1
Goat, edible offal of	*0.1
Goat meat	*0.1
Sheep, edible offal of	3
Sheep meat	0.2
Albendazole sulfoxide see Albendazole	
Aldicarb Sum of aldicarb, its sulfoxide and its sulfone, expressed as aldicarb	
Citrus fruits	0.05
Cotton seed	*0.05
Edible offal (mammalian)	*0.01
Meat (mammalian)	*0.01
Milks	*0.01
Sugar cane	*0.02
Aldoxycarb Sum of aldoxycarb and its sulfone, expressed as aldoxycarb	
Cattle, edible offal of	0.2
Cattle meat	*0.02
Eggs	0.1
Milks	*0.02
Poultry, edible offal of	0.2
Poultry meat	*0.02
Wheat	*0.02
Aliphatic alcohol ethoxylates Aliphatic alcohol ethoxylates	
Cattle, edible offal of	*0.1
Cattle meat	*0.1
Cattle milk	1
Altrenogest Altrenogest	
Pig meat	*0.005
Pig, edible offal of	0.005
Aluminium phosphide see Phosphine	

Ametoctradin <i>Commodities of plant origin:</i> Ametoctradin <i>Commodities of animal origin:</i> Sum of ametoctradin and 6-(7-amino-5-ethyl [1,2,4] triazolo [1,5-a]pyrimidin-6-yl) hexanoic acid	
Edible offal (mammalian)	*0.02
Eggs	*0.02
Grapes	3
Meat (mammalian)	*0.02
Milks	*0.02
Poultry, edible offal of	*0.02
Poultry meat	*0.02
Ametryn Ametryn	
Cotton seed	0.05
Edible offal (mammalian)	*0.05
Meat (mammalian)	*0.05
Milks	*0.05
Pineapple	*0.05
Pome fruits	0.1
Sugar cane	0.05
Aminoethoxyvinylglycine Aminoethoxyvinylglycine	
Apple	0.1
Stone fruits [except cherries]	0.2
Walnuts	*0.05
Aminopyralid <i>Commodities of plant origin:</i> Sum of aminopyralid and conjugates, expressed as aminopyralid <i>Commodities of animal origin:</i> Aminopyralid	
Cereal grains	0.1
Edible offal (mammalian) [except kidney]	0.02
Eggs	*0.01
Kidney (mammalian)	0.3
Meat (mammalian)	*0.01
Milks	*0.01
Poultry, edible offal of	*0.01
Poultry meat	*0.01
Wheat bran, unprocessed	0.3
Amitraz Sum of amitraz and N-(2,4-dimethylphenyl)-n'-methylformamidine, expressed as N-(2,4-dimethylphenyl)-N'-methylformamidine	
Apple	0.5
Cotton seed	*0.1
Cotton seed oil, crude	1
Edible offal (mammalian)	0.5
Meat (mammalian)	0.1
Milks	0.1
Stone fruits [except cherries]	0.5
Amitrole Amitrole	
Avocado	*0.01
Banana	*0.01
Blueberries	T*0.01
Cereal grains	*0.01
Citrus fruits	*0.01
Edible offal (mammalian)	*0.01
Grapes	*0.01

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Hops, dry	*0.01	Sweet corn (corn-on-the-cob)	*0.1
Meat (mammalian)	*0.01		
Milks	*0.01	Avermectin B1	
Oilseed	*0.01	see Abamectin	
Papaya (pawpaw)	*0.01		
Passionfruit	*0.01	Avilamycin	
Pecan	*0.01	Inhibitory substance, identified as avilamycin	
Pineapple	*0.01	Poultry, edible offal of	*0.05
Pome fruits	*0.01	Poultry meat	*0.05
Potato	*0.05		
Pulses	*0.01	Azaconazole	
Stone fruits	*0.02	Azaconazole	
Sugar cane	*0.01	Mushrooms	0.1
Amoxycillin		Azamethiphos	
Inhibitory substance, identified as amoxycillin		Azamethiphos	
Cattle milk	*0.01	Cereal grains	0.1
Edible offal (mammalian)	*0.01	Edible offal (mammalian)	*0.05
Eggs	*0.01	Eggs	*0.05
Meat (mammalian)	*0.01	Meat (mammalian)	*0.05
Poultry, edible offal of	*0.01	Milks	*0.05
Poultry meat	*0.01	Poultry, edible offal of	*0.05
Sheep milk	*0.01	Poultry meat	*0.05
		Wheat bran, unprocessed	0.5
Ampicillin			
Inhibitory substance, identified as ampicillin		Azaperone	
Cattle milk	*0.01	Azaperone	
Horse, edible offal of	*0.01	Pig, edible offal of	0.2
Horse meat	*0.01	Pig meat	0.2
Amprolium		Azimsulfuron	
Amprolium		Azimsulfuron	
Eggs	4	Edible offal (mammalian)	*0.02
Poultry, edible offal of	1	Eggs	*0.02
Poultry meat	0.5	Meat (mammalian)	*0.02
		Milks	*0.02
Apramycin		Poultry, edible offal of	*0.02
Apramycin		Poultry meat	*0.02
Edible offal (mammalian)	2	Rice	*0.02
Meat (mammalian)	*0.05		
Poultry, edible offal of	1	Azinphos-methyl	
Poultry meat	*0.05	Azinphos-methyl	
		Blueberries	1
Asulam		Citrus fruits	2
Asulam		Edible offal (mammalian)	*0.05
Apple	*0.1	Grapes	2
Edible offal (mammalian)	*0.1	Kiwifruit	2
Hops, dry	*0.1	Litchi	2
Meat (mammalian)	*0.1	Macadamia nuts	*0.01
Milks	*0.1	Meat (mammalian)	*0.05
Poppy seed	*0.1	Milks	*0.05
Potato	0.4	Oilseed	*0.05
Sugar cane	*0.1	Pome fruits	2
		Raspberries, red, black	1
Atrazine		Stone fruits	2
Atrazine		Strawberry	1
Edible offal (mammalian)	T*0.1		
Lupin (dry)	*0.02	Azoxystrobin	
Maize	*0.1	Azoxystrobin	
Meat (mammalian)	T*0.01	Almonds	*0.01
Milks	T*0.01	Anise myrtle leaves	T100
Potato	*0.01	Avocado	1
Rape seed (canola)	*0.02	Banana	T0.5
Sorghum	*0.1	Barley	*0.02
Sugar cane	*0.1	Beans [except broad and soya bean]	2

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Bergamot	T50	Tea, green, black	T20
Blackberries	5	Tomato	T1
Blueberries	5	Tree nuts [except almonds]	2
Boysenberry	5	Turmeric, root	T0.1
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicac	0.7	Wheat	*0.02
Brassica leafy vegetables [except mizuna]	2	Bacitracin	
Bulb vegetables [except fennel, bulb; onion, bulb]	2	Inhibitory substance, identified as bacitracin	
Burnet, Salad	T50	Chicken, edible offal of	*0.5
Carrot	0.2	Chicken fat	*0.5
Chervil	T50	Chicken meat	*0.5
Chick-pea (dry)	T0.5	Eggs	*0.5
Citrus fruits	10	Milks	*0.5
Cloudberry	T5	Benalaxyl	
Coriander (leaves, stem, roots)	T50	Benalaxyl	
Coriander, seed	T50	Fruiting vegetables, cucurbits	0.2
Cotton seed	*0.01	Garlic	0.1
Cranberry	0.5	Grapes	0.5
Dewberries (including loganberry)	T5	Lettuce, head	*0.01
Dill, seed	T50	Lettuce, leaf	*0.01
Dried grapes	5	Onion, bulb	0.1
Edible offal (mammalian)	*0.01	Shallot	T0.5
Eggs	*0.01	Spring onion	T0.1
Fennel, seed	T50	Bendiocarb	
Fennel, bulb	T0.1	<i>Commodities of plant origin:</i> Unconjugated bendiocarb	
Fruiting vegetables, cucurbits	1	<i>Commodities of animal origin:</i> Sum of conjugated and unconjugated Bendiocarb, 2,2-dimethyl-1,3- benzodioxol-4-ol and N-hydroxymethylbendiocarb, expressed as Bendiocarb	
Galangal, Greater	T0.1	Banana	*0.02
Grapes	2	Cattle, edible offal of	0.2
Herbs [except as otherwise listed under this chemical]	T50	Cattle meat	0.1
Horseradish	0.5	Eggs	0.05
Kaffir lime leaves	T50	Milks	0.1
Lemon grass	T50	Poultry, edible offal of	0.1
Lemon myrtle leaves	T100	Poultry meat	0.05
Lemon verbena (dry leaves)	T50	Benfluralin	
Lentil (dry)	T0.5	Benfluralin	
Lettuce, head	15	Lettuce, head	T*0.05
Lettuce, leaf	15	Lettuce, leaf	T*0.05
Maize	T*0.01	Benomyl	
Mango	0.5	see Carbendazim	
Meat (mammalian)	*0.01	Bensulfuron-methyl	
Mexican tarragon	T50	Bensulfuron-methyl	
Milks	0.005	Rice	*0.02
Mizuna	T50	Rice bran, processed	*0.05
Olives	T2	Bensulide	
Passionfruit	0.5	Bensulide	
Peanut	0.05	Fruiting vegetables, cucurbits	*0.1
Peanut oil, crude	0.1	Bentazone	
Peas (pods and succulent, immature seeds)	2	Bentazone	
Peppers	3	Beans [except broad bean and soya bean]	*0.1
Poppy seed	*0.02	Broad bean (green pods and immature seeds)	*0.1
Potato	0.05	Edible offal (mammalian)	*0.05
Poultry, edible offal of	*0.01	Eggs	*0.05
Poultry meat	*0.01		
Radish	0.5		
Raspberries, red, black	5		
Riberries	T10		
Rice	T7		
Rose and dianthus (edible flowers)	T50		
Spices	*0.1		
Stone fruits	1.5		
Strawberry	10		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Garden pea (shelled)	T*0.05	Peas	T0.5
Meat (mammalian)	*0.05	Peppers	T0.5
Milks	*0.05	Plums (including prunes)	0.5
Onion, bulb	T0.1	Pome fruits	2
Peanut	*0.1	Raspberries, red, black	T7
Podded pea (young pods) (snow and sugar snap)	T0.05	Sinkwa or Sinkwa towel gourd	T0.5
Poultry, edible offal of	*0.05	Squash, Summer	T0.5
Poultry meat	*0.05	Strawberry	T2
Pulses	*0.01	Tomato	T1
Rice	*0.03	Watermelon	T0.3
Sweet corn (corn-on-the-cob)	*0.1	Yard-long bean (pods)	T1
Benzocaine Benzocaine		Bifenthrin Bifenthrin	
Abalone	*0.05	Apple	*0.05
Finfish	*0.05	Avocado	T0.1
Benzofenap Sum of benzofenap, benzofenap-OH and Benzofenap-red, expressed as benzofenap		Banana	0.1
Rice	*0.01	Blackberries	T3
Benzyladenine Benzyladenine		Blueberries	T3
Apple	0.2	Brassica (cole or cabbage) vegetables, Head cabbages, Flower head brassicas [except Cabbages, Head]	T1
Pear	T0.2	Cabbages, Head	T7
Pistachio nut	T*0.05	Cereal grains	*0.02
Benzyl G penicillin Inhibitory substance, identified as benzyl G penicillin		Cherries	T1
Edible offal (mammalian)	*0.06	Chervil	T0.5
Meat (mammalian)	*0.06	Chia	T0.2
Milks	*0.0015	Cloudberry	T3
Betacyfluthrin see Cyfluthrin		Citrus fruits	*0.05
Bifenazate Sum of bifenazate and bifenazate diazene (diazene-carboxylic acid, 2-(4-methoxy-[1,1'-biphenyl-3-yl] 1-methylethyl ester), expressed as bifenazate		Common bean (pods and/or immature seeds)	T1
Almonds	0.1	Cotton seed	0.1
Apricot	0.5	Cucumber	T0.5
Bitter melon	T0.5	Dewberries (including boysenberry and loganberry)	T3
Blackberries	T7	Edible offal (mammalian)	0.5
Cherries	2.5	Eggs	*0.05
Cloudberry	T7	Field pea (dry)	T*0.01
Cranberry	1.5	Fruiting vegetables, cucurbits [except cucumber]	0.1
Cucumber	T0.5	Fruiting vegetables, other than cucurbits	0.5
Dewberries (including boysenberry and loganberry)	T7	Galangal, rhizomes	T10
Dried grapes	T2	Ginger, root	T*0.01
Edible offal (mammalian)	*0.01	Gooseberry	T3
Egg plant	T0.1	Grapes	*0.01
Grapes [except wine grapes]	T1	Herbs	T0.5
Hops, dry	T3	Kaffir lime leaves	T10
Lettuce, head	T20	Leafy vegetables [except chervil; mizuna; rucola (rocket)]	T2
Lettuce, leaf	T20	Lemon balm	T10
Meat (mammalian) (in the fat)	*0.01	Lemon grass	T10
Melons, except watermelon	T0.3	Lemon verbena	T10
Milks	*0.01	Lupin (dry)	T*0.02
Nectarine	0.5	Meat (mammalian) (in the fat)	2
Papaya (pawpaw)	T0.5	Milks	0.5
Peach	2	Mizuna	T0.5
		Olives	T0.5
		Pear	0.5
		Peas (pods and succulent, immature seeds)	*0.01
		Pineapple	T*0.01
		Poppy seed	*0.02
		Poultry, edible offal of	*0.05
		Poultry meat (in the fat)	*0.05

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Pulses [except field pea (dry) and lupin (dry)]	*0.02	Raspberries, red, black	T10
Rape seed (canola)	*0.02	Root and tuber vegetables	1
Raspberries, red, black	T3	Silvanberries	T10
Rucola (rocket)	T0.5	Stone fruits [except cherries]	1.7
Stone fruits [except cherries]	1	Strawberry	10
Strawberry	1		
Sugar cane	*0.01	Brodifacoum	
Sweet potato	*0.05	Brodifacoum	
Taro	T*0.05	Cereal grains	T*0.00002
Tea, green, black	5	Edible offal (mammalian)	T*0.00005
Turmeric, root	T10	Meat (mammalian)	T*0.00005
		Pulses	T*0.00002
		Sugar cane	*0.0005
Bioresmethrin			
Bioresmethrin		Bromacil	
Mango	T0.5	Bromacil	
Bitertanol		Asparagus	*0.04
Bitertanol		Citrus fruits	*0.04
Beans [except broad bean and soya bean]	0.5	Edible offal (mammalian)	*0.04
Edible offal (mammalian)	3	Meat (mammalian)	*0.04
Eggs	*0.01	Milks	*0.04
Meat (mammalian) (in the fat)	0.3	Pineapple	*0.04
Milks	0.2		
Poultry, edible offal of	*0.01	Bromoxynil	
Poultry meat	*0.01	Bromoxynil	
Strawberry	*0.05	Cereal grains	*0.2
		Edible offal (mammalian)	T3
		Eggs	*0.02
		Garlic	T0.1
		Grapes	*0.01
		Linseed	*0.02
		Meat (mammalian) (in the fat)	T1
		Milks	T0.1
		Poultry, edible offal of	*0.02
		Poultry meat	*0.02
		Sugar cane	*0.02
Boscalid			
Commodities of plant origin: Boscalid		Bupirimate	
Commodities of animal origin: Sum of boscalid, 2-chloro-N-(4'-chloro-5-hydroxybiphenyl-2-yl)nicotinamide and the glucuronide conjugate of 2-chloro-N-(4'-chloro-5-hydroxybiphenyl-2-yl)nicotinamide, expressed as boscalid equivalents		Bupirimate	
All other foods	0.5	Apple	1
Blackberries	T10	Egg plant	T1
Blueberries	T15	Fruiting vegetables, cucurbits	1
Boysenberry	T10	Peppers	0.7
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	2	Strawberry	1
Bulb vegetables [except onion, bulb]	T5		
Celery	T15	Buprofezin	
Cherries	T3	Buprofezin	
Chervil	T30	Celery	T5
Cloudberry	T10	Chervil	T50
Coriander (leaves, roots and stems)	T30	Citrus fruits	2
Dewberries (including loganberry and youngberry) [except boysenberry]	T10	Coriander (leaves, stem, roots)	T50
Dried grapes	15	Cotton seed	T1
Fruiting vegetables, cucurbits	0.5	Cotton seed oil, crude	T0.3
Fruiting vegetables, other than cucurbits	1	Custard apple	0.1
Edible offal (mammalian)	0.3	Dried grapes (currants, raisins and sultanas)	1
Grapes	4	Edible offal (mammalian)	*0.05
Herbs	T30	Fruiting vegetables, cucurbits	T2
Leafy vegetables	30	Fruiting vegetables, other than cucurbits	T2
Legume vegetables	3	Grapes	0.3
Meat (mammalian) (in the fat)	0.3	Herbs	T50
Milk fats	0.7	Lettuce, leaf	T10
Milks	0.1	Mango	0.2
Onion, bulb	T1	Meat (mammalian) (in the fat)	*0.05
Pistachio nut	T2		
Pome fruits	2		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Milks	*0.01	Poultry meat	*0.02
Mizuna	T50	Stone fruits	15
Olives	T0.5	Strawberry	10
Olive oil, crude	T2	Tree nuts [except almonds]	3
Passionfruit	2		
Pear	0.2		
Persimmon, Japanese	1		
Rucola (rocket)	T50		
Stone fruits [except apricot; peach]	1.9		
Tree tomato	T1		
Butafenacil		Carbaryl	
Butafenacil		Carbaryl	
Cereal grains [except rice]	*0.02	Apricot	10
Edible offal (mammalian)	*0.02	Asparagus	10
Eggs	*0.01	Avocado	10
Grapes	T*0.02	Banana (in the pulp)	5
Meat (mammalian)	*0.01	Barley	15
Milks	*0.01	Blackberries	10
Pome fruits	T*0.02	Blueberries	7
Poultry, edible offal of	*0.02	Brazilian cherry (grumichama)	5
Poultry meat	*0.01	Carambola	5
Stone fruits	T*0.02	Cassava	T0.1
		Cereal grains [except barley; sorghum]	5
		Cherries	5
		Citrus fruits	7
		Cotton seed	3
		Cranberry	3
		Custard apple	5
		Dewberries (including boysenberry and loganberry)	10
		Edible offal (mammalian)	T0.2
		Eggs	T0.2
		Elephant apple	5
		Feijoa	5
		Fruiting vegetables, cucurbits	3
		Galangal, rhizomes (fresh)	T5
		Granadilla	5
		Grapes	5
		Guava	5
		Jaboticaba	5
		Jackfruit	5
		Jambu	5
		Kiwifruit	10
		Leafy vegetables	10
		Litchi	5
		Longan	5
		Mango	5
		Meat (mammalian)	T0.2
		Milks	T*0.05
		Nectarine	10
		Okra	10
		Olives	10
		Olives, processed	1
		Papaya (pawpaw)	5
		Passionfruit	5
		Peach	10
		Plums (including prunes)	5
		Pome fruits	5
		Potato	0.2
		Poultry, edible offal of	T5
		Poultry meat	T0.5
		Rambutan	5
		Raspberries, red, black	10
		Sapodilla	5
		Sapote, black	5
		Sapote, green	5
		Sapote, mammey	5
		Sapote, white	5
		Sorghum	10
		Strawberry	7
		Sugar cane	T*0.05
Butroxydim			
Butroxydim			
Edible offal (mammalian)	*0.01		
Eggs	*0.01		
Legume vegetables	*0.01		
Meat (mammalian)	*0.01		
Milks	*0.01		
Oilseed	*0.01		
Poultry, edible offal of	*0.01		
Poultry meat	*0.01		
Pulses	*0.01		
Cadusafos			
Cadusafos			
Banana	*0.01		
Citrus fruits	*0.01		
Ginger, root	0.1		
Sugar cane	*0.01		
Tomato	*0.01		
Captan			
Captan			
Almonds	0.3		
Berries and other small fruits [except blueberries; grapes; strawberry]	T30		
Blueberries	20		
Chick-pea (dry)	T0.1		
Cucumber	T5		
Dried grapes	15		
Edible offal (mammalian)	*0.05		
Eggs	*0.02		
Grapes	10		
Lentil (dry)	T0.1		
Lettuce, leaf	T7		
Meat (mammalian)	*0.05		
Milks	*0.01		
Peppers, Chili	T7		
Peppers, Sweet	T7		
Pitaya (dragon fruit)	T20		
Pome fruits	10		
Poultry, edible offal of	*0.02		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Sunflower seed	1	Wheat	0.2
Sweet corn (corn-on-the-cob)	1		
Tree nuts	1	Carbon disulphide	
Tree nuts (whole in shell)	10	Carbon disulfide	
Turmeric, root (fresh)	T5	Cereal grains	10
Vegetables [except as otherwise listed under this chemical]	5	Pulses	T10
Wheat bran, unprocessed	T20		
Carbendazim		Carbonyl sulphide	
Sum of carbendazim and 2-aminobenzimidazole, expressed as carbendazim		Carbonyl sulphide	
Apple	0.2	Cereal grains	T0.2
Apricot	2	Pulses	T0.2
Banana	T1	Rape seed (canola)	T0.2
Berries and other small fruits [except grapes]	T5		
Cherries	20	Carbosulfan	
Chives	*0.1	see Carbofuran	
Citron	0.7		
Edible offal (mammalian)	0.2	Carboxin	
Eggs	*0.1	Carboxin	
Garlic	T0.2	Cereal grains	0.1
Ginger, root	T10		
Grapefruit	0.2	Carfentrazone-ethyl	
Grapes	0.3	Carfentrazone-ethyl	
Lemon	0.7	Assorted tropical and sub-tropical fruits – edible peel	*0.05
Lime	0.7	Assorted tropical and sub-tropical fruits – inedible peel	*0.05
Macadamia nuts	0.1	Berries and other small fruits [except grapes]	T*0.05
Mandarins	0.7	Cereal grains	*0.05
Meat (mammalian)	0.2	Citrus fruits	*0.05
Milks	*0.1	Cotton seed	T*0.05
Mineola	0.7	Edible offal (mammalian)	*0.05
Mushrooms	T5	Eggs	*0.05
Nectarine	0.2	Grapes	*0.05
Onion, bulb	T*0.2	Hops, dry	*0.05
Oranges	0.2	Meat (mammalian)	*0.05
Peach	0.2	Milks	*0.025
Pear	0.2	Pome fruits	*0.05
Peppers	*0.1	Potato	*0.05
Peppers, Chili (dry)	20	Poultry, edible offal of	*0.05
Poultry, edible offal of	*0.1	Poultry meat	*0.05
Poultry meat	*0.1	Stone fruits	*0.05
Pulses	0.5	Tree nuts	*0.05
Shaddock (pomelo)	0.2		
Spices	*0.1	Ceftiofur	
Sugar cane	T0.1	Desfuroylceftiofur	
Tangelo [except mineola]	0.2	Cattle, edible offal of	2
Tangors	0.7	Cattle fat	0.5
Tomato	0.5	Cattle meat	0.1
		Cattle milk	0.1
Carbofuran			
Sum of carbofuran and 3-hydroxycarbofuran, expressed as carbofuran		Cefuroxime	
Barley	0.2	Inhibitory substance, identified as cefuroxime	
Cotton seed	0.1	Cattle, edible offal of	*0.1
Edible offal (mammalian)	*0.05	Cattle meat	*0.1
Eggs	*0.05	Cattle milk	*0.1
Garlic	T0.1		
Meat (mammalian)	*0.05	Cephalonium	
Milks	*0.05	Inhibitory substance, identified as cephalonium	
Poultry, edible offal of	*0.05	Cattle, edible offal of	*0.1
Poultry meat	*0.05	Cattle meat	*0.1
Rice	0.2	Cattle milk	*0.02
Sugar cane	*0.1		
Sunflower seed	0.1		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Cephapirin		Turnip, Garden	T0.05
Cephapirin and des-acetylcephapirin, expressed as cephapirin		Chlorfenapyr	
Cattle, edible offal of	*0.02	Chlorfenapyr	
Cattle meat	*0.02	Brassica (cole or cabbage) vegetables,	0.5
Cattle milk	*0.01	Head cabbages, Flowerhead	
Chinomethionat		brassicas	
see Oxythioquinox		Brassica leafy vegetables [except	T3
Chlorantraniliprole		chinese cabbage]	
<i>Plant commodities and animal commodities other than milk: Chlorantraniliprole</i>		Chinese cabbage	3
<i>Milk: Sum of chlorantraniliprole, 3-bromo-N-[4-chloro-2-(hydroxymethyl)-6-[(methylamino)carbonyl]phenyl]-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide, and 3-bromo-N-[4-chloro-2-(hydroxymethyl)-6-[[[(hydroxymethyl)amino]carbonyl]phenyl]-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide, expressed as chlorantraniliprole</i>		Cotton seed	0.5
Adzuki bean (dry)	T0.5	Edible offal (mammalian)	*0.05
All other foods	*0.01	Eggs	*0.01
Almonds	T0.05	Meat (mammalian) (in the fat)	0.05
Brassica (cole or cabbage) vegetables,	0.5	Milks	*0.01
Head cabbages, Flowerhead		Mizuna	T3
brassicas		Onion, Welsh	T1
Celery	5	Peach	1
Chick-pea (dry)	0.07	Pome fruits	0.5
Cotton seed	0.3	Poultry, edible of	*0.01
Coriander (leaves, stem, roots)	T20	Poultry meat (in the fat)	*0.01
Cranberry	1	Rucola (rocket)	T5
Dried fruits	2	Shallot	T1
Edible offal (mammalian) [except liver]	*0.01	Spring onion	T1
Eggs	0.03	Chlorfenvinphos	
Fruiting vegetables, cucurbits	0.2	Chlorfenvinphos, sum of E and Z isomers	
Fruiting vegetables, other than	0.3	Broccoli	T0.05
cucurbits [except peppers, chili and		Brussels sprouts	T0.05
sweet corn (corn-on-the-cob)]		Cabbages, head	T0.05
Grapes [except table grapes]	0.3	Carrot	T0.4
Herbs	T20	Cattle, edible offal of	T*0.1
Leafy vegetables [except lettuce, head;	15	Cattle meat (in the fat)	T0.2
rucola]		Cattle milk (in the fat)	T0.2
Legume vegetables	1	Cauliflower	T0.1
Lettuce, head	3	Celery	T0.4
Liver (mammalian)	0.02	Cotton seed	T0.05
Meat (mammalian) (in the fat)	0.02	Deer meat (in the fat)	0.2
Mexican tarragon	T20	Egg plant	T0.05
Milk fats	0.1	Goat, edible offal of	T*0.1
Milks	*0.01	Goat meat (in the fat)	T0.2
Mung bean (dry)	0.7	Horseradish	T0.1
Peppers, Chili	1	Leek	T0.05
Pistachio nut	T0.05	Maize	T0.05
Pome fruits	0.3	Mushrooms	T0.05
Potato	*0.01	Onion, bulb	T0.05
Poultry, edible offal of	*0.01	Peanut	T0.05
Poultry meat (in the fat)	*0.01	Potato	T0.05
Radish	T0.05	Radish	T0.1
Rhubarb	5	Rice	T0.05
Rucola (rocket)	T20	Sheep, edible offal of	T*0.1
Soya bean (dry)	0.07	Sheep meat (in the fat)	T0.2
Stone fruits	1	Swede	T0.05
Strawberry	T0.5	Sweet potato	T0.05
Swede	T0.05	Tomato	T0.1
Sweet corn (corn-on-the-cob)	*0.01	Turnip, garden	T0.05
Table grapes	1.2	Wheat	T0.05
Chlorfluazuron		Chlorfluazuron	
		Chlorfluazuron	
		Cattle, edible offal of	0.1
		Cattle meat (in the fat)	1
		Cattle milk	0.1
		Cotton seed	0.1
		Cotton seed oil, crude	0.1
		Cotton seed oil, edible	*0.05

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Eggs	0.2	Meat (mammalian) (in the fat)	2
Poultry, edible offal of	0.1	Milks	0.05
Poultry meat (in the fat)	1	Nectarine	7
Chlorhexidine		Onion, bulb	10
Chlorhexidine		Onion, Welsh	T10
Milks	0.05	Papaya (pawpaw)	10
Sheep, edible offal of	*0.5	Peach	30
Sheep fat	*0.5	Peanut	0.2
Sheep meat	*0.5	Peas (pods and succulent, immature seeds)	10
Chloridazon		Persimmon, American	T5
Chloridazon		Persimmon, Japanese	T5
Beetroot	*0.05	Plums (including prunes)	10
Chlormequat		Potato	0.1
Chlormequat cation		Poultry, edible offal of	*0.05
Barley	T2	Poultry meat	*0.05
Dried grapes	0.75	Pulses	3
Edible offal (mammalian)	0.5	Rice	T*0.1
Eggs	0.1	Shallot	T10
Grapes	0.75	Spring onion	T10
Meat (mammalian)	0.2	Sunflower seed	T*0.01
Milks	0.5	Tomato	10
Poultry, edible offal of	0.1	Tree tomato	T10
Poultry meat	*0.05	Turmeric root	T7
Wheat	5	Vegetables [except asparagus; Brussels sprouts; carrot; celery; egg plant; fennel bulb; fruiting vegetables; cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spring onion; tomato]	T7
Chloropicrin		Wasabi	T7
Chloropicrin		Chlorpropham	
Cereal grains	*0.1	Chlorpropham	
Chlorothalonil		Garlic	*0.05
<i>Commodities of plant origin: Chlorothalonil</i>		Onion, bulb	*0.05
<i>Commodities of animal origin: 4-hydroxy-2,5,6-trichloroisophthalonitrile metabolite, expressed as chlorothalonil</i>		Potato	30
Almonds	T0.1	Chlorpyrifos	
Apricot	7	Chlorpyrifos	
Asparagus	T*0.1	Asparagus	T0.5
Banana	3	Avocado	0.5
Berries and other small fruits [except blackcurrant and grapes]	T10	Banana	T0.5
Brussels sprouts	7	Blackberries	0.5
Carrot	7	Blueberries	*0.01
Celery	10	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	T0.5
Cherries	10	Cassava	T*0.02
Coriander (leaves, stem, roots)	T20	Celery	T5
Currant, black	10	Cereal grains [except sorghum]	T0.1
Edible offal (mammalian)	7	Cherries	1
Egg plant	T10	Citrus fruits	T0.5
Fennel, bulb	5	Coffee beans	T0.5
Fennel, leaf	5	Cotton seed	0.05
Fennel, seed	5	Cotton seed oil, crude	0.2
Fruiting vegetables, cucurbits	5	Cranberry	1
Galangal, Greater	T7	Dried fruits	T2
Galangal, Lesser	T7	Edible offal (mammalian)	T0.1
Garlic	10	Eggs	T*0.01
Grapes	10	Ginger, root	*0.02
Herbs [except fennel, leaf]	T20	Grapes	T1
Leafy vegetables [except lettuce]	T100	Kiwifruit	2
Leek	T10	Leek	T5
Lettuce, head	T10	Mango	*0.05
Lettuce, leaf	T10		
Mango	T1		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Meat (mammalian) (in the fat)	T0.5	Pig meat	0.1
Milks (in the fat)	T0.2	Poultry, edible offal of	0.6
Oilseed [except cotton seed and peanut]	T*0.05	Poultry meat	0.1
Olives	T*0.05	Chlorthal-dimethyl Chlorthal-dimethyl	
Parsley	0.05	Eggs	*0.05
Passionfruit	*0.05	Edible offal (mammalian)	*0.05
Peanut	0.05	Meat (mammalian)	*0.05
Peppers, Chili (dry)	20	Lettuce, head	2
Peppers, Sweet	T1	Lettuce, leaf	2
Persimmon, American	T1	Milks	*0.05
Persimmon, Japanese	T1	Parsley	T2
Pineapple	T0.5	Poultry, edible offal of	*0.05
Pitaya (dragon fruit)	T*0.05	Poultry meat	*0.05
Pome fruits	T0.5	Vegetables [except as otherwise listed under this chemical]	5
Potato	0.05	Clavulanic acid Clavulanic acid	
Poultry, edible offal of	T0.1	Cattle, edible offal of	*0.01
Poultry meat (in the fat)	T0.1	Cattle meat	*0.01
Sorghum	T3	Cattle milk	*0.01
Spices	5	Clethodim see Sethoxydim	
Star apple	T*0.05	Clodinafop-propargyl Clodinafop-propargyl	
Stone fruits [except cherries]	T1	Barley	T*0.02
Strawberry	0.3	Edible offal (mammalian)	*0.05
Sugar cane	T0.1	Eggs	*0.05
Swede	T0.3	Meat (mammalian)	*0.05
Sweet potato	T0.05	Milks	*0.05
Taro	0.05	Poultry, edible offal of	*0.05
Tea, green, black	2	Poultry meat	*0.05
Tomato	T0.5	Wheat	*0.05
Tree nuts	T0.05	Clodinafop acid (R)-2-[4-(5-chloro-3-fluoro-2-pyridinyloxy) phenoxy] propanoic acid	
Vegetables [except asparagus; brassica vegetables; cassava; celery; leek; peppers, chili (dry); peppers, Sweet; potato; swede; sweet potato; taro and tomato]	T*0.01	Barley	T*0.02
Chlorpyrifos-methyl Chlorpyrifos-methyl		Edible offal (mammalian)	*0.1
Cereal grains [except rice]	10	Eggs	*0.1
Cotton seed	*0.01	Meat (mammalian)	*0.1
Edible offal (mammalian)	*0.05	Milks	*0.1
Eggs	*0.05	Poultry, edible offal of	*0.1
Lupin (dry)	10	Poultry meat	*0.1
Meat (mammalian) (in the fat)	*0.05	Wheat	*0.1
Milks (in the fat)	*0.05	Clofentezine Clofentezine	
Poultry, edible offal of	*0.05	Almonds	T0.5
Poultry meat (in the fat)	*0.05	Banana	*0.01
Rice	0.1	Edible offal (mammalian)	T*0.05
Wheat bran, unprocessed	20	Grapes	1
Wheat germ	30	Hops, dry	*0.2
Chlorsulfuron Chlorsulfuron		Meat (mammalian)	T*0.05
Cereal grains	*0.05	Milks	T*0.05
Edible offal (mammalian)	*0.05	Pome fruits	0.1
Meat (mammalian)	*0.05	Stone fruits	0.1
Milks	*0.05	Tomato	T1
Chlortetracycline Inhibitory substance, identified as chlortetracycline			
Cattle kidney	0.6		
Cattle liver	0.3		
Cattle meat	0.1		
Eggs	0.2		
Pig kidney	0.6		
Pig liver	0.3		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Clomazone Clomazone			
Beans [except broad bean and soya beans]	*0.05	Maize	*0.01
Common beans (pod and/or immature seeds)	T*0.05	Meat (mammalian)	*0.02
Fruiting vegetables, cucurbits	*0.05	Milks	*0.01
Poppy seed	*0.05	Olives	T0.5
Potato	*0.05	Persimmon, American	T2
Rice	*0.01	Persimmon, Japanese	T2
		Pome fruits	T2
		Popcorn	*0.01
		Poultry, edible offal of	*0.02
		Poultry meat	*0.02
		Rape seed (canola)	*0.01
		Sorghum	*0.01
		Soursop	T2
		Soya bean (dry)	T0.02
		Stone fruits [except cherries]	T3
		Sugar apple	T2
		Sugar cane	0.1
		Sunflower seed	*0.01
		Sweet corn (corn-on-the-cob)	0.2
		Wine grapes	*0.02
Clopyralid Clopyralid		Cloxacillin Inhibitory substance, identified as Cloxacillin	
Cauliflower	T0.2	Cattle milk	*0.01
Cereal grains	2		
Edible offal (mammalian) [except kidney]	0.5		
Hops, dry	2		
Kidney of cattle, goats, pigs and sheep	5		
Meat (mammalian)	0.1		
Milks	0.05		
Poppy seed	T0.5		
Rape seed (canola)	0.5		
Cloquintocet-mexyl Sum of cloquintocet mexyl and 5-chloro-8-quinolinoxyacetic acid, expressed as cloquintocet mexyl		Coumaphos Sum of coumaphos and its oxygen analogue, expressed as coumaphos	
Barley	*0.1	Cattle fat	*0.02
Edible offal (mammalian)	*0.1	Cattle kidney	*0.02
Eggs	*0.1	Cattle liver	*0.02
Meat (mammalian)	*0.1	Cattle milk	*0.01
Milks	*0.1	Cattle milk fat	0.1
Poppy seed	T*0.02	Cattle muscle	*0.02
Poultry, edible offal of	*0.1		
Poultry meat	*0.1		
Rye	*0.1		
Triticale	*0.1		
Wheat	*0.1		
Clorsulon Clorsulon		Cyanamide Cyanamide	
Cattle, edible offal of	*0.1	Apple	*0.02
Cattle meat	*0.1	Blueberries	*0.05
Cattle milk	1.5	Grapes	*0.05
		Kiwifruit	*0.1
		Pear, Oriental (nashi)	*0.1
		Plums (including prunes)	*0.02
Closantel Closantel		Cyanazine Cyanazine	
Sheep, edible offal of	5	Bulb vegetables	*0.02
Sheep meat	2	Cereal grains	*0.01
		Leek	0.05
		Peas	0.02
		Podded pea (young pods) (snow and sugar snap)	0.05
		Potato	0.02
		Pulses	*0.01
		Sweet corn (corn-on-the-cob)	*0.02
Clothianidin Clothianidin			
Apricot	T2		
Banana	*0.02		
Cherimoya	T2		
Cherries	T5		
Cotton seed	*0.02		
Cranberry	0.01		
Custard apple	T2		
Dried grapes	10		
Edible offal (mammalian)	*0.02		
Eggs	*0.02		
Grapes [except wine grapes]	3		
Llama	T2		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Cyantraniliprole <i>Commodities of plant origin:</i> Cyantraniliprole <i>Commodities of animal origin for enforcement:</i> Cyantraniliprole <i>Commodities of animal origin for dietary exposure assessment:</i> Sum of cyantraniliprole and 2-[3-bromo-1-(3-chloropyridin-2-yl)-1H-pyrazol-5-yl]-3,8-dimethyl-4-oxo-3,4-dihydroquinazoline-6-carbonitrile (IN-J9Z38), 2-[3-bromo-1-(3-chloropyridin-2-yl)-1H-pyrazol-5-yl]-8-methyl-4-oxo-3,4-dihydroquinazoline-6-carbonitrile (IN-MLA84), 3-bromo-1-(3-chloropyridin-2-yl)-N-[4-cyano-2-[(hydroxymethyl)carbamoyl]-6-methylphenyl]-1H-pyrazole-5-carboxamide (IN-MYX98) and 3-bromo-1-(3-chloropyridin-2-yl)-N-[4-cyano-2-(hydroxymethyl)-6-(methylcarbamoyl)phenyl]-1H-pyrazole-5-carboxamide (IN-N7B69), expressed as cyantraniliprole		Carambola	T0.1
		Cereal grains	2
		Chia	T0.5
		Citrus fruits	0.2
		Cotton seed	0.01
		Cotton seed oil, crude	0.02
		Custard apple	T0.1
		Edible offal (mammalian)	*0.01
		Egg plant	T0.2
		Eggs	*0.01
		Grapes	1
		Legume vegetables	0.5
		Lemon aspen	T1
		Litchi	T0.1
		Macadamia nuts	0.05
		Mango	T0.1
		Mammalian fats [except milk fats]	0.5
		Meat (mammalian)	0.02
		Milks	0.1
		Okra	T0.2
		Papaya (pawpaw)	T0.2
		Pecan	T0.05
		Peppers, Sweet	T0.2
		Persimmon, American	T0.1
		Persimmon, Japanese	T0.1
		Poultry, edible offal of	*0.01
		Poultry meat (in the fat)	*0.01
		Pulses	0.5
		Rape seed (canola)	*0.05
		Stone fruits	0.3
		Tomato	0.2
		Wheat bran, unprocessed	5
Cyclanilide Sum of cyclanilide and its methyl ester, expressed as cyclanilide		Cyhalofop-butyl Sum of cyhalofop-butyl, cyhalofop and metabolites expressed as cyhalofop-butyl	
All other foods	0.05	Edible offal (mammalian)	*0.05
Bulb vegetables [except onion, bulb]	7	Eggs	*0.05
Cotton seed	*0.01	Meat (mammalian) (in the fat)	*0.05
Edible offal (mammalian)	*0.01	Milks	*0.05
Eggs	*0.01	Poultry, edible offal of	*0.05
Fruiting vegetables, cucurbits	0.5	Poultry meat	*0.05
Fruiting vegetables, other than cucurbits	2	Rice	*0.01
Meat (mammalian) (in the fat)	*0.01	Cyhalothrin Cyhalothrin, sum of isomers	
Milk fats	*0.01	Barley	0.2
Milks	*0.01	Beetroot	*0.01
Onion, bulb	0.05	Berries and other small fruits	0.2
Potato	0.05	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	0.1
Poultry, edible offal of	*0.01	Cereal grains [except barley; sorghum; wheat]	*0.01
Poultry meat (in the fat)	*0.01	Chard	T0.5
Cyflufenamid Cyflufenamid		Citrus fruits	*0.01
Dried grapes (currants, raisins and sultanas)	0.5	Coriander (leaves, stem, roots)	T1
Edible offal (mammalian)	*0.01	Cotton seed	*0.02
Eggs	*0.01	Cucumber	T0.05
Fruiting vegetables, cucurbits	0.1	Edible offal (mammalian)	*0.02
Grapes	0.15	Eggs	*0.02
Meat (mammalian) (in the fat)	*0.01	Garlic	*0.05
Milks	*0.01	Legume vegetables	0.1
Poultry, edible offal of	*0.01	Meat (mammalian) (in the fat)	0.5
Poultry meat (in the fat)	*0.01	Milks (in the fat)	0.5
Cyfluthrin Cyfluthrin, sum of isomers		Onion, bulb	*0.05
Avocado	0.1		
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	0.5		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Onion, Welsh	T0.05	Peas	1
Parsley	T1	Peppers, Chili	1
Potato	*0.01	Pig, edible offal of	*0.05
Poultry, edible offal of	*0.02	Pig meat (in the fat)	*0.05
Poultry meat	*0.02	Persimmon, American	T2
Pulses [except soya bean (dry)]	0.2	Persimmon, Japanese	T2
Radish	*0.01	Pome fruits	1
Rape seed (canola)	0.02	Poppy seed	T*0.01
Shallot	T0.05	Potato	*0.01
Sorghum	0.5	Poultry, edible offal of	*0.05
Soya bean (dry)	*0.02	Poultry meat (in the fat)	*0.05
Spring onion	T0.05	Radish	T0.05
Stone fruits	0.5	Rape seed (canola)	0.2
Sunflower seed	*0.01	Rape seed oil, edible	0.2
Tea, green, black	1	Shallot	T0.5
Tomato	0.02	Sheep, edible offal of	0.05
Wheat	*0.05	Sheep meat (in the fat)	0.5
Cypermethrin		Soya bean (dry)	0.05
Cypermethrin, sum of isomers		Soya bean oil, crude	0.1
Adzuki bean (dry)	T0.05	Spring onion	T0.5
All other foods	*0.01	Stone fruits	1
Asparagus	0.5	Sunflower seed	0.1
Avocado	T0.2	Sunflower seed oil, crude	0.1
Beetroot	T0.1	Sweet corn (corn-on-the-cob)	0.05
Berries and other small fruits [except grapes]	0.5	Tea, green, black	0.5
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	1	Tomato	0.5
Broad bean (dry) (fava bean)	0.05	Wheat	0.2
Cattle, edible offal of	0.05	Cyproconazole	
Cattle meat (in the fat)	0.5	Cyproconazole, sum of isomers	
Celery	T1	Barley	*0.02
Cereal grains [except wheat]	1	Chick-pea (dry)	T*0.01
Chick-pea (dry)	0.2	Edible offal (mammalian)	1
Common bean (dry) (navy bean)	0.05	Eggs	*0.01
Coriander (leaves, stem, roots)	T5	Lentil (dry)	T*0.01
Coriander, seed	T1	Meat (mammalian)	0.03
Cotton seed	0.2	Milks	*0.01
Cotton seed oil, crude	*0.02	Peanut	0.02
Cucumber	T0.3	Potato	*0.02
Deer meat (in the fat)	T0.5	Poultry, edible offal of	*0.01
Durian	1	Poultry meat	*0.01
Eggs	0.05	Wheat	*0.02
Field pea (dry)	0.05	Cyprodinil	
Goat, edible offal of	0.05	Cyprodinil	
Goat meat (in the fat)	0.5	Blackberries	10
Grapes	T0.05	Blueberries	3
Herbs	T5	Boysenberry	10
Horse, edible offal of	*0.05	Cloudberry	T5
Horse meat (in the fat)	*0.05	Common bean (pods and/or immature seeds)	0.7
Leafy vegetables [except lettuce head]	T5	Cucumber	0.5
Leek	T0.5	Dewberries (including boysenberry and loganberry)	T5
Lemon balm	T5	Dried grapes (currants, raisins and sultanas)	5
Lettuce, head	2	Dried stone fruits	0.05
Linola oil, edible	0.1	Edible offal (mammalian)	*0.01
Linola seed	0.1	Egg plant	T0.2
Linseed	0.5	Grapes	2
Longan	1	Leafy vegetables	10
Lupin (dry)	*0.01	Meat (mammalian)	*0.01
Milks (in the fat)	1	Melons, except watermelon	T0.2
Mung bean (dry)	0.05	Milks	*0.01
Olives	T*0.05	Onion, bulb	0.2
Onion, bulb	*0.01	Peas (pods and succulent, immature)	0.5
Onion, Welsh	T0.5		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

seeds)					
Peppers, Sweet	0.7				
Pistachio nut	T0.1				
Pome fruits	0.05				
Raspberries, red, black	10				
Stone fruits	2				
Strawberry	5				
Tomato	T1				
Cyromazine					
Cyromazine					
Cattle, edible offal of	0.05				
Cattle meat	0.05				
Eggs	0.2				
Goat, edible offal of	0.2				
Goat meat	0.2				
Milks	*0.01				
Mushrooms	10				
Pig, edible offal of	0.05				
Pig meat	0.05				
Poultry, edible offal of	0.1				
Poultry meat	0.05				
Sheep, edible offal of	0.2				
Sheep meat	0.2				
2,4-D					
2,4-D					
Cereal grains	0.2				
Citrus fruits	5				
Edible offal (mammalian)	2				
Eggs	*0.05				
Grapes	T*0.05				
Legume vegetables	*0.05				
Lupin (dry)	*0.05				
Meat (mammalian)	0.2				
Milks	*0.05				
Oilseed	*0.05				
Pear	*0.05				
Potato	0.1				
Poultry, edible offal of	*0.05				
Poultry meat	*0.05				
Pulses	*0.05				
Sugar cane	5				
Daminozide					
Daminozide					
Edible offal (mammalian)	0.2				
Eggs	0.2				
Meat (mammalian)	0.2				
Milks	*0.05				
Peach	30				
Peanut	20				
Pome fruits	30				
Poultry, edible offal of	0.2				
Poultry meat	0.2				
2,4-DB					
2,4-DB					
Cereal grains	*0.02				
Edible offal (mammalian)	0.2				
Eggs	*0.05				
Meat (mammalian)	0.2				
Milks	*0.05				
Poultry, edible offal of	*0.05				
Poultry meat	*0.05				
Deltamethrin					
Deltamethrin					
Brassica (cole or cabbage) vegetables,	*0.05				
Head cabbages, Flowerhead					
brassicas					
Cattle, edible offal of	0.1				
Cattle meat (in the fat)	0.5				
Cereal grains	2				
Eggs	*0.01				
Fruiting vegetables, other than	0.1				
cucurbits					
Goat, edible offal of	0.1				
Goat meat (in the fat)	0.2				
Legume vegetables	0.1				
Milks	0.05				
Oilseed	0.1				
Pig, edible offal of	*0.01				
Pig meat (in the fat)	0.1				
Poultry, edible offal of	*0.01				
Poultry meat (in the fat)	*0.01				
Pulses	0.1				
Sheep, edible offal of	0.1				
Sheep meat (in the fat)	0.2				
Sweet corn (kernels)	0.1				
Tea, green, black	5				
Wheat bran, unprocessed	5				
Wheat germ	3				
Derquantel					
Derquantel					
Sheep fat	0.0002				
Sheep kidney	0.0002				
Sheep liver	0.0002				
Sheep muscle	0.0002				
Dexamethasone and Dexamethasone					
trimethylacetate					
Dexamethasone					
Cattle, edible offal of	0.1				
Cattle meat	0.1				
Cattle milk	*0.05				
Horse, edible offal of	0.1				
Horse meat	0.1				
Pig, edible offal of	0.1				
Pig meat	0.1				
Diafenthiuron					
Sum of diafenthiuron; N-[2,6-bis(1-methylethyl)-4-phenoxyphe-nyl]-N'-(1,1-dimethylethyl)urea; and N-[2,6-bis(1-methylethyl)-4-phenoxyphe-nyl]-N'-(1,1-dimethylethyl)carbodiimide, expressed as diafenthiuron					
Cotton seed	0.2				
Edible offal (mammalian)	*0.02				
Eggs	*0.02				
Meat (mammalian) (in the fat)	*0.02				
Milks	*0.02				
Peanut	T0.1				
Poultry, edible offal of	*0.02				
Poultry meat (in the fat)	*0.02				
Diazinon					
Diazinon					
Cereal grains	0.1				
Citrus fruits	0.7				

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Coriander (leaves, stem, roots)	*0.05
Coriander, seed	*0.05
Edible offal (mammalian)	0.7
Eggs	*0.05
Fruit [except as otherwise listed under this chemical]	0.5
Kiwifruit	0.5
Meat (mammalian) (in the fat)	0.7
Milks (in the fat)	0.5
Olive oil, crude	2
Parsley	*0.05
Peach	0.7
Poultry, edible offal of	*0.05
Poultry meat	*0.05
Shallot	T0.5
Spring onion	T0.5
Sugar cane	0.5
Sweet corn (corn-on-the-cob)	0.7
Tree nuts	0.1
Vegetable oils, crude [except olive oil, virgin]	0.1
Vegetables	0.7
Dicamba Dicamba	
Cereal grains	*0.05
Edible offal (mammalian)	0.05
Eggs	*0.05
Meat (mammalian)	0.05
Milks	0.1
Poultry, edible offal of	*0.05
Poultry meat	*0.05
Sugar cane	0.1
Sugar cane molasses	2
Dicamba Sum of dicamba, 3,6-dichloro-5-hydroxy-2-methoxybenzoic acid and 3,6-dichloro-2-hydroxybenzoic acid, expressed as dicamba	
Soya bean	10
Dichlobenil Dichlobenil	
Blueberries	T1
Citrus fruits	0.1
Currants, black, red, white	T1
Gooseberry	T1
Grapes	0.1
Pome fruits	0.1
Raspberries, red, black	T1
Stone fruits	0.1
Tomato	0.1
Dichlofluanid Dichlofluanid	
Berries and other small fruits [except grapes and strawberry]	T50
Grapes	0.5
Peanut	*0.02
Strawberry	10
Tomato	1

1,3-dichloropropene 1,3-dichloropropene	
Grapes	0.018
Dichlorprop-P Sum of dichlorprop acid, its esters and conjugates, hydrolysed to dichlorprop acid, and expressed as dichlorprop acid	
Citrus fruits	0.2
Edible offal (mammalian)	*0.05
Eggs	*0.02
Meat (mammalian)	*0.02
Milks	*0.01
Poultry, edible offal of	*0.05
Poultry meat	*0.02
Dichlorvos Dichlorvos	
Cacao beans	5
Cereal grains	5
Coffee beans	2
Edible offal (mammalian)	0.05
Eggs	0.05
Fruit	0.1
Lentil (dry)	2
Lettuce, head	1
Lettuce, leaf	1
Meat (mammalian)	0.05
Milks	0.02
Mushrooms	0.5
Peanut	2
Poultry, edible offal of	0.05
Poultry meat	0.05
Rape seed (canola)	T0.1
Rice bran, unprocessed	10
Soya bean (dry)	2
Tomato	0.5
Tree nuts	2
Vegetables [except as otherwise listed under this chemical]	0.5
Wheat bran, unprocessed	10
Wheat germ	10
Diclofop-methyl Diclofop-methyl	
Cereal grains	0.1
Edible offal (mammalian)	*0.05
Eggs	*0.05
Lupin (dry)	0.1
Meat (mammalian)	*0.05
Milks	*0.05
Oilseed	0.1
Peas	0.1
Poppy seed	0.1
Poultry, edible offal of	*0.05
Poultry meat	*0.05
Dicloran Dicloran	
Beans [except broad bean and soya bean]	20
Berries and other small fruits [except grapes]	20
Broad bean (green pods and immature seeds)	20

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Carrot	15	Pome fruits	0.3
Grapes	10	Potato	*0.02
Lettuce, head	20	Poultry meat	*0.05
Lettuce, leaf	20	Poultry, edible offal of	*0.05
Onion, bulb	20	Spinach	T3
Stone fruits	15	Tomato	0.5
Sweet potato	20		
Tomato	20		
Dicofol		Diflubenzuron	
Sum of dicofol and 2,2,2- trichloro-1-(4-chlorophenyl)-1-(2-chlorophenyl)ethanol, expressed as dicofol		Diflubenzuron	
Almonds	5	Cattle, edible offal of	*0.02
Cotton seed	0.1	Cattle milk	0.05
Cucumber	2	Cereal grains	T2
Fruit [except strawberry]	5	Mushrooms	0.1
Gherkin	2	Sheep kidney	0.05
Hops, dry	5	Sheep liver	0.05
Strawberry	1	Sheep meat (in the fat)	0.05
Tea, green, black	5	Sheep milk	0.05
Tomato	1	Wheat bran, unprocessed	T5
Vegetables [except as otherwise listed under this chemical]	5		
Dicyclanil		Diflufenican	
Sum of dicyclanil and its triaminopyridyl metabolite expressed as dicyclanil		Diflufenican	
Sheep fat	0.3	Barley	0.05
Sheep kidney	0.3	Edible offal (mammalian)	0.1
Sheep liver	0.3	Eggs	*0.02
Sheep meat	0.3	Grapes	*0.002
		Meat (mammalian)	0.01
		Milks	0.01
		Oats	0.05
		Peas	0.05
		Poultry, edible offal of	*0.02
		Poultry meat	*0.02
		Pulses	0.05
		Rye	0.05
		Triticale	0.05
		Wheat	0.02
Didecyldimethylammonium chloride		Dimethenamid-P	
Didecyldimethylammonium chloride		Sum of dimethenamid-P and its (R)-isomer	
Assorted tropical and sub-tropical fruits – inedible peel	20	Common bean (pods and/or immature seeds)	*0.02
		Edible offal (mammalian)	*0.01
		Eggs	*0.01
		Maize	*0.02
		Meat (mammalian)	*0.01
		Milks	*0.01
		Peas	*0.02
		Poppy seed	*0.01
		Poultry, edible offal of	*0.01
		Poultry meat	*0.01
		Pulses	*0.02
		Pumpkins	*0.02
		Rape seed (canola)	T*0.01
		Sweet corn (corn-on-the-cob)	*0.02
Dieldrin		Dimethipin	
see Aldrin and Dieldrin		Dimethipin	
		Cotton seed	0.5
		Cotton seed oil, crude	*0.1
		Cotton seed oil, refined	*0.1
		Edible offal (mammalian)	*0.01
		Eggs	*0.02
		Meat (mammalian)	*0.01
		Milks	*0.01
		Poultry, edible offal of	*0.01
Difenoconazole			
Difenoconazole			
Asparagus	*0.05		
Avocado	0.5		
Banana	*0.02		
Beetroot	T0.5		
Carrot	0.2		
Cereal grains	*0.01		
Celeriac	T0.5		
Celery	T5		
Chard (silverbeet)	T3		
Chicory leaves (green and red cultivars)	T3		
Chives	2		
Dried grapes	6		
Edible offal (mammalian)	*0.05		
Eggs	*0.05		
Endive	T3		
Grapes	4		
Macadamia nuts	*0.01		
Meat (mammalian)	*0.05		
Milks	*0.01		
Papaya (pawpaw)	1		
Parsley	T15		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Poultry meat	*0.01	Sweet potato	0.1
Dimethirimol		Tomato	0.02
Dimethirimol		Turnip, garden	*0.2
Fruiting vegetables, cucurbits	1	Watermelon	T5
Dimethoate		Wheat bran, processed	T1
Sum of dimethoate and omethoate, expressed as dimethoate <i>see also</i> Omethoate		Dimethomorph	
Abiu	5	Sum of E and Z isomers of dimethomorph	
Artichoke, globe	T1	Beetroot	T0.1
Asparagus	0.02	Edible offal (mammalian)	*0.01
Assorted tropical and sub-tropical fruits – inedible peel [except avocado; mango]	5	Fruiting vegetables, cucurbits	0.5
Avocado	3	Grapes	2
Banana passionfruit	5	Leafy vegetables [except lettuce head]	T10
Bearberry	T5	Leek	0.5
Beetroot	T*0.1	Lettuce, head	0.3
Bilberry	T5	Meat (mammalian)	*0.01
Bilberry, bog	T5	Milks	*0.01
Bilberry, red	T5	Mizuna	T10
Blackberries	T5	Onion, bulb	0.05
Blueberries	T5	Onion, Welsh	2
Boysenberry	0.02	Parsley	T2
Broccoli	T0.3	Peas	1
Cabbages, head	T0.2	Poppy seed	*0.02
Cactus fruit	5	Potato	*0.02
Carrot	T0.3	Radish	T0.1
Cauliflower	T0.3	Shallot	T0.5
Celery	T0.5	Spring onion	2
Cereal grains	T0.05	Dinitolmide	
Cherries	T0.2	Sum of dinitolmide and its metabolite 3-amino-5-nitro-o-toluamide, expressed as dinitolmide equivalents	
Citrus fruits	5	Poultry, edible offal of	6
Cranberry	T5	Poultry fats	2
Edible offal (mammalian)	0.1	Poultry meat	3
Egg plant	T0.02	Dinitro-o-toluamide	
Eggs	*0.05	<i>see</i> Dinitolmide	
Elderberries	0.02		
Grapes	T*0.1	Dinotefuran	
Legume vegetables	T2	Sum of dinotefuran and its metabolites DN, 1-methyl-3-(tetrahydro-3-furylmethyl)guanidine and UF, 1-methyl-3-(tetrahydro-3-furylmethyl)urea expressed as dinotefuran	
Mango	1	Grapes	0.9
Meat (mammalian)	*0.05	Diphenylamine	
Melons, except watermelon	T5	Diphenylamine	
Milks	*0.05	Apple	10
Oilseed [except peanut]	T0.1	Edible offal (mammalian) [except liver]	*0.01
Olive oil, refined	T0.1	Eggs	0.05
Onion, bulb	0.7	Liver of cattle, goats, pigs and sheep	0.05
Parsnip	T0.3	Meat (mammalian) (in the fat)	*0.01
Peanut	T*0.05	Milks (in the fat)	*0.01
Peppers, Chili	T5	Pear	7
Peppers, Sweet	0.7	Poultry, edible offal of	*0.01
Potato	0.1	Poultry meat (in the fat)	*0.01
Poultry, edible offal of	*0.05	Diquat	
Poultry meat	*0.05	Diquat cation	
Pulses	T0.5	Anise myrtle leaves	T0.5
Radish	T3	Barley	5
Raspberries, red, black	T5	Beans [except broad bean and soya bean]	1
Rhubarb	0.7	Broad bean (green pods and immature	1
Rollinia	5		
Santols	5		
Squash, summer (including zucchini)	0.7		
Stone fruits [except cherries]	T*0.02		
Strawberry	0.02		
Sweet corn (corn-on-the-cob)	T0.3		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

seeds)		Beetroot	1
Edible offal (mammalian)	*0.05	Berries and other small fruits [except strawberry]	T10
Eggs	*0.01	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	2
Fruit	*0.05	Broad bean (green pods and immature seeds)	2
Hops, dry	T0.2	Bulb vegetables [except garlic and onion, bulb]	T10
Lemon myrtle leaves	T0.5	Carrot	1
Linseed	*0.01	Celery	5
Maize	0.1	Cereal grains	0.5
Meat (mammalian)	*0.05	Citrus fruits	0.2
Milks	*0.01	Coconut	5
Native pepper (Tasmania lanceolata) leaves	T0.5	Coffee beans	5
Oats	5	Common bean (pods and/or immature seeds)	2
Oilseed [except linseed and poppy seed]	5	Cotton seed	10
Onion, bulb	0.1	Custard apple	5
Peas	0.1	Edible offal (mammalian)	2
Poppy seed	*0.01	Eggs	*0.5
Potato	0.2	Fig	3
Poultry, edible offal of	*0.05	Fruiting vegetables, cucurbits	2
Poultry meat	*0.05	Fruiting vegetables, other than cucurbits [except roselle]	3
Pulses	1	Garlic	4
Rice	5	Herbs [except parsley]	T5
Rice, polished	1	Hops	T10
Rye	2	Leafy vegetables	5
Sorghum	2	Litchi	5
Sugar beet	0.1	Macadamia nuts	*0.2
Sugar cane	*0.05	Mango	7
Tea, green, black	T0.5	Meat (mammalian)	*0.5
Tree nuts	*0.05	Milks	*0.2
Triticale	2	Onion, bulb	4
Vegetable oils, crude	1	Papaya (pawpaw)	5
Vegetables [except beans; broad bean; onion, bulb; peas; potato; pulses; sugar beet]	*0.05	Parsley	5
Wheat	2	Parsnip	T1
Disulfoton		Passionfruit (including Granadilla)	3
Sum of disulfoton and demeton-S and their sulfoxides and sulfones, expressed as disulfoton		Peanut	0.2
Cotton seed	0.5	Peas (pods and succulent, immature seeds)	2
Edible offal (mammalian)	0.02	Persimmon, Japanese	3
Eggs	*0.02	Pistachio nut	T3
Hops, dry	0.5	Pome fruits	3
Meat (mammalian)	0.02	Pomegranate	3
Milks	0.01	Poppy seed	*0.2
Potato	0.5	Potato	1
Poultry, edible offal of	*0.02	Poultry meat	*0.5
Poultry meat	*0.02	Poultry, edible offal of	*0.5
Vegetables	0.5	Pulses	0.5
Dithianon		Radish	T1
Dithianon		Rhubarb	2
Fruit	2	Roselle (rosella)	5
Dithiocarbamates		Stone fruits	3
Total dithiocarbamates, determined as carbon disulphide evolved during acid digestion and expressed as milligrams of carbon disulphide per kilogram of food		Strawberry	5
Almonds	3	Sunflower seed	T*0.05
Asparagus	T1	Swede	T1
Avocado	7	Tree tomato	T5
Banana	2	Turnip, garden	T1
Beans [except broad bean and soya bean]	2	Walnuts	T*0.2
		Wasabi	T2

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Diuron			
Sum of diuron and 3,4- dichloroaniline, expressed as diuron			
Asparagus	2	Dill, seed	T0.05
Cereal grains	0.1	Edible offal (mammalian)	0.02
Cotton seed oil, crude	0.5	Egg plant	T0.1
Edible offal (mammalian)	3	Fennel, seed	T0.05
Fruit	0.5	Grapes	*0.002
Meat (mammalian)	0.1	Herbs	T0.05
Milks	0.1	Kaffir lime leaves	T0.05
Oilseed	0.5	Leafy vegetables [except lettuce head; lettuce leaf; mizuna]	T0.5
Pulses	*0.05	Lemon grass	T0.05
Sugar cane	0.2	Lemon verbena (fresh weight)	T0.05
		Lettuce, head	0.2
		Lettuce, leaf	0.2
		Meat (mammalian)(in the fat)	0.01
		Milks	*0.001
		Milk fats	0.01
		Mizuna	T0.5
		Parsnip	T0.05
		Peppers, Sweet	0.01
		Pulses	*0.01
		Radish	T0.05
		Rape seed (canola)	*0.01
		Strawberry	T0.1
		Swede	T0.05
		Sweet corn (corn-on-the-cob)	*0.002
		Tomato	0.01
		Turnip, garden	T0.05
Dodine		Endosulfan	
Dodine		Sum of A- and B- endosulfan and endosulfan sulphate	
Pome fruits	5	Assorted tropical and sub-tropical fruits – inedible peel	2
Stone fruits	*0.05	Broccoli	1
		Cabbages, head	1
		Cauliflower	1
		Cereal grains	0.1
		Citrus fruits	0.3
		Edible offal (mammalian)	0.2
		Eggs	0.02
		Fruiting vegetables, cucurbits	1
		Fruiting vegetables, other than cucurbits	1
		Meat (mammalian) (in the fat)	0.2
		Milks	0.02
		Oilseed	1
		Pome fruits	1
		Poultry, edible offal of	*0.01
		Poultry meat (in the fat)	0.05
		Pulses	*0.1
		Root and tuber vegetables	0.5
		Stalk and stem vegetables	1
		Strawberry	T0.5
		Tea, green, black	T30
		Tree nuts	0.05
Doramectin		Endothal	
Doramectin		Endothal	
Cattle, edible offal of	0.1	Cotton seed	0.1
Cattle fat	0.1	Potato	0.1
Cattle meat	0.01		
Cattle milk	0.05		
Pig kidney	0.03		
Pig liver	0.05		
Pig meat (in the fat)	0.1		
Sheep, edible offal of	0.05		
Sheep fat	0.1		
Sheep meat	0.02		
2,2-DPA		Enilconazole	
2,2-dichloropropionic acid		see Imazalil	
Avocado	*0.1		
Banana	*0.1		
Cereal grains	*0.1		
Citrus fruits	*0.1		
Cotton seed	*0.1		
Currants, black, red, white	15		
Edible offal (mammalian)	0.2		
Grapes	3		
Meat (mammalian)	0.2		
Milks	*0.1		
Papaya (pawpaw)	*0.1		
Pecan	*0.1		
Pineapple	*0.1		
Pome fruits	*0.1		
Stone fruits	1		
Sugar cane	*0.1		
Sunflower seed	*0.1		
Vegetables	*0.1		
EDC			
see Ethylene dichloride			
Emamectin			
Sum of emamectin B1a and emamectin B1b			
Beetroot	T0.05		
Bergamot	T0.05		
Burnet, salad	T0.05		
Celery	T0.2		
Coriander (leaves, stem, roots)	T0.05		
Coriander, seed	T0.05		
Cotton seed	0.005		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Epoxiconazole Epoxiconazole			
Avocado	0.5	Olives	T5
Banana	1	Oranges, sweet, sour	2
Cereal grains	0.05	Peach	0.5
Edible offal (mammalian)	0.05	Pineapple	2
Eggs	*0.01	Poultry, edible offal of	*0.2
Meat (mammalian)	*0.01	Poultry meat	*0.1
Milks	*0.005	Sugar cane	0.5
Poultry, edible offal of	*0.01	Sugar cane molasses	7
Poultry meat (in the fat)	*0.01	Tomato	2
Wheat bran, unprocessed	0.3	Walnuts	T5
Wheat germ	0.2	Wheat	T1
Eprinomectin Eprinomectin B1a		Ethion Ethion	
Cattle, edible offal of	2	Cattle, edible offal of	2.5
Cattle fat	0.5	Cattle meat (in the fat)	2.5
Cattle milk	0.03	Citrus fruits	1
Cattle meat	0.1	Cotton seed	0.1
Deer, edible offal of	2	Cotton seed oil, crude	0.05
Deer meat	0.1	Grapes	2
EPTC EPTC		Milks (in the fat)	0.5
Cereal grains	*0.04	Pome fruits	1
Edible offal (mammalian)	*0.1	Stone fruits	1
Eggs	*0.01	Tea, green, black	5
Meat (mammalian)	*0.1	Ethofumesate Ethofumesate	
Milks	*0.1	Beetroot	0.1
Oilseed	0.1	Bulb vegetables	*0.1
Poultry, edible offal of	*0.05	Chard (silver beet)	1
Poultry meat	*0.05	Edible offal (mammalian)	0.5
Vegetables	*0.04	Meat (mammalian) (in the fat)	0.5
Erythromycin Inhibitory substance, identified as erythromycin		Milks (in the fat)	0.2
Edible offal (mammalian)	*0.3	Poppy seed	*0.02
Meat (mammalian)	*0.3	Spinach	T1
Milks	*0.04	Sugar beet	0.1
Poultry, edible offal of	*0.3	Ethopabate Ethopabate	
Poultry meat	*0.3	Poultry, edible offal of	15
Esfenvalerate see Fenvalerate		Poultry meat	5
Ethephon Ethephon		Ethoprophos Ethoprophos	
Apple	1	Banana	*0.05
Banana	T*0.05	Cereal grains	*0.005
Barley	1	Custard apple	*0.02
Cherries	15	Litchi	*0.02
Cotton seed	2	Potato	*0.02
Cotton seed oil, crude	*0.1	Sugar cane	*0.1
Currant, black	1	Sweet potato	*0.02
Edible offal (mammalian)	0.2	Tomato	*0.01
Eggs	*0.2	Ethoxyquin Ethoxyquin	
Grapes	10	Apple	3
Kiwifruit	0.1	Pear	3
Macadamia nuts	*0.1	Ethoxysulfuron <i>Commodities of plant origin:</i> Ethoxysulfuron <i>Commodities of animal origin:</i> 2-amino-4, 6-dimethoxypyrimidine, expressed as ethoxysulfuron	
Mandarins	2	Edible offal (mammalian)	*0.05
Mango	T*0.02	Meat (mammalian)	*0.05
Meat (mammalian)	0.1	Milks	*0.01
Milks	0.1		
Nectarine	0.01		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Sugar cane	*0.01	lettuce, leaf]	0.2
Ethyl formate		Lettuce, head	0.2
Ethyl formate		Lettuce, leaf	0.2
Dried fruits	1	Meat (mammalian)	*0.05
Ethylene dichloride (EDC)		Milks	*0.005
1,2-dichloroethane		Mushrooms	0.1
Cereal grains	*0.1	Onion, bulb	*0.05
Etoxazole		Peanut	*0.05
Etoxazole		Pineapple	*0.05
Banana	0.2	Poultry, edible offal of	*0.05
Cherries	1	Poultry meat	*0.05
Chervil	T1	Root and tuber vegetables	0.2
Citrus fruits	0.5	Strawberry	0.2
Coriander (leaves, stem, roots)	T1	Sugar cane	*0.05
Cotton seed	0.2	Tomato	0.5
Custard apple	T0.1	Fenarimol	
Dried grapes	1.5	Fenarimol	
Edible offal (mammalian)	*0.01	Berries and other small fruits [except grapes]	T0.1
Eggs	*0.01	Cherries	1
Fruiting vegetables, other than cucurbits	0.05	Fruiting vegetables, cucurbits	0.2
Fruiting vegetables, cucurbits	T0.1	Grapes	0.1
Grapes	0.5	Pome fruits	0.2
Herbs	T1	Fenbendazole	
Ivy gourd	T0.1	Fenbendazole	
Meat (mammalian) (in the fat)	*0.02	Cattle, edible offal of	*0.1
Milks	*0.01	Cattle meat	*0.1
Mizuna	T1	Goat, edible offal of	0.5
Papaya	T0.1	Goat meat	0.5
Podded pea (young pods) (snow and sugar snap)	T0.1	Milks	0.1
Pointed gourd	T0.1	Sheep, edible offal of	0.5
Pome fruits	0.2	Sheep meat	0.5
Poultry, edible offal of	*0.01	Fenbuconazole	
Poultry meat (in the fat)	*0.02	Fenbuconazole	
Rucola (Rocket)	T1	Banana	0.5
Stone fruits [except cherries]	0.3	Blueberries	0.3
Etridiazole		Edible offal (mammalian)	0.05
Etridiazole		Eggs	*0.01
Beetroot	*0.02	Meat (mammalian)	*0.01
Cotton seed	*0.02	Milks	*0.01
Peanut	*0.02	Nectarine	0.5
Vegetables [except as otherwise listed under this chemical]	0.2	Poultry, edible offal of	*0.01
Fenamiphos		Poultry meat	*0.01
Sum of fenamiphos, its sulfoxide and sulfone, expressed as fenamiphos		Stone fruits [except nectarine]	1
Aloe vera	1	Wheat	*0.01
Banana	*0.05	Fenbutatin oxide	
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	*0.05	Bis[tris(2-methyl-2-phenylpropyl)tin]-oxide	
Celery	*0.05	Assorted tropical and sub-tropical fruits – inedible peel	5
Citrus fruits	*0.05	Berries and other small fruits [except table grapes]	1
Edible offal (mammalian)	*0.05	Cherries	6
Eggs	*0.05	Citrus fruits	5
Fruiting vegetables, cucurbits	*0.05	Citrus peel	30
Ginger, root	*0.05	Dried grapes	T10
Grapes	*0.05	Fig	T10
Leafy vegetables [except lettuce, head;	*0.05	Grapes [except wine grapes]	T3
		Hops, dry	20
		Nectarine	3
		Peach	3
		Pome fruits	3

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Tomato	T2	Fenoxaprop-ethyl Sum of fenoxaprop-ethyl (all isomers) and 2-(4-(6-chloro-2-benzoxazolyloxy)phenoxy)-propanoate and 6-chloro-2,3-dihydrobenzoxazol-2-one, expressed as fenoxaprop-ethyl	
Fenhexamid Fenhexamid		Fenoxycarb Fenoxycarb	
Blackberries	T20	Barley	*0.01
Blueberries	5	Chick-pea (dry)	*0.01
Chervil	T15	Edible offal (mammalian)	0.2
Cloudberry	T20	Eggs	*0.02
Coriander (leaves, stem, roots)	T15	Meat (mammalian)	0.05
Cucumber	T10	Milks	0.02
Dewberries (including boysenberry, loganberry and youngberry)	T20	Poultry, edible offal of	*0.1
Dried grapes	20	Poultry meat	*0.01
Edible offal (mammalian)	2	Rice	T*0.02
Grapes	10	Rye	*0.01
Herbs	T15	Triticale	*0.01
Kiwifruit	15	Wheat	*0.01
Lettuce, head	T50	Fenpropathrin Fenpropathrin	
Lettuce, leaf	T50	Cherries	5
Meat (mammalian) (in the fat)	*0.05	Citrus fruits	2
Milks	*0.01	Grapes	5
Mizuna	T15	Tea, green, black	2
Peas (pods and succulent, immature seeds)	T5	Fenpyroximate Fenpyroximate	
Peppers	T30	Apple	0.3
Raspberries, red, black	T20	Citrus fruits	0.6
Rucola (rocket)	T15	Pear	0.3
Stone fruits [except plums]	10	Strawberry	1
Strawberry	10	Fenthion Sum of fenthion, its oxygen analogue, and their sulfoxides and sulfones, expressed as fenthion	
Tomato	T2	Apricot	T0.2
Fenitrothion Fenitrothion		Assorted tropical and sub-tropical fruits – inedible peel	5
Apple	0.5	Cattle, edible offal of	1
Cabbages, head	0.5	Cattle meat	1
Cacao beans	0.1	Cherries	T0.4
Cereal grains	10	Citrus fruits	T0.7
Cherries	0.5	Eggs	*0.05
Edible offal (mammalian)	*0.05	Grapes	T0.2
Eggs	*0.05	Melons, except watermelon	T3
Fruit [except as otherwise listed under this chemical]	0.1	Milks	T0.2
Grapes	0.5	Nectarine	T0.25
Lettuce, head	0.5	Olive oil, crude	T0.5
Lettuce, leaf	0.5	Olives	T0.2
Meat (mammalian)	T*0.05	Peach	T0.2
Milks (in the fat)	T*0.05	Peppers, Chili	T7
Oilseeds	T0.1	Peppers, Sweet	T0.5
Poultry, edible offal of	*0.05	Persimmon, Japanese	T0.3
Poultry meat	*0.05	Pig, edible offal of	0.5
Pulses [except soya bean (dry)]	T0.1	Pig meat	0.5
Rice, polished	0.1	Plums	T0.25
Soya bean (dry)	0.3		
Sugar cane	0.02		
Tea, green, black	0.5		
Tomato	0.5		
Tree nuts	0.1		
Vegetables [except as otherwise listed under this chemical]	0.1		
Wheat bran, unprocessed	20		
Wheat germ	20		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Pome fruits	T0.25	Head cabbages, Flowerhead	
Poultry, edible offal of	*0.05	brassicas	
Poultry meat	*0.05	Burnet, salad	T0.1
Sheep, edible offal of	0.2	Celery	T0.3
Sheep meat	0.2	Chervil	T0.1
Watermelon	T3	Citrus fruits	T*0.01
Fentin		Coriander (leaves, stem, roots)	T0.1
Fentin hydroxide, excluding inorganic tin and Di- and Mono-phenyltin		Coriander, seed	T0.1
Cacao beans	*0.1	Cotton seed	*0.01
Carrot	0.2	Cotton seed oil, crude	*0.01
Celeriac	0.1	Custard apple	T0.05
Celery	1	Dill, seed	T0.1
Coffee beans	*0.1	Edible offal (mammalian)	0.02
Peanut	*0.05	Eggs	0.02
Pecan	*0.05	Fennel, seed	T0.1
Potato	0.1	Ginger, root	*0.01
Rice	*0.1	Grapes [except wine grapes]	T*0.01
Sugar beet	0.2	Herbs	T0.1
Fenvalerate		Honey	0.01
Fenvalerate, sum of isomers		Kaffir lime leaves	T0.1
Berries and other small fruits	1	Lemon grass	T0.1
Brassica (cole or cabbage) vegetables,	1	Lemon verbena (fresh weight)	T0.1
Head cabbages, Flowerhead		Lettuce, head	T0.1
brassicas		Lettuce, leaf	T0.1
Brassica leafy vegetables	1	Meat (mammalian) (in the fat)	0.1
Cereal grains	2	Milks	0.01
Celery	2	Mizuna	T0.1
Dried grapes	0.5	Mushrooms	0.02
Edible offal (mammalian)	0.05	Peanut	T*0.01
Eggs	0.02	Peanut oil, crude	T*0.01
Grapes	0.1	Pecan	T*0.01
Legume vegetables	0.5	Peppers, Chili	*0.005
Meat (mammalian) (in the fat)	1	Peppers, Sweet	T0.1
Milks	0.2	Pome fruits	T*0.01
Oilseed [except peanut]	0.5	Poppy seed	*0.01
Peanut	T0.1	Potato	*0.01
Pome fruits	1	Poultry, edible offal of	*0.01
Poultry, edible offal of	*0.02	Poultry meat (in the fat)	0.02
Poultry meat (in the fat)	0.05	Rape seed (canola)	*0.01
Pulses	0.5	Rice	*0.005
Stone fruits	1	Rucola (rocket)	T0.1
Sweet corn (corn-on-the-cob)	0.05	Sorghum	0.01
Tea, green, black	0.05	Stone fruits	0.01
Tomato	0.2	Sugar cane	*0.01
Wheat bran, unprocessed	5	Sunflower seed	*0.01
Fipronil		Swede	0.1
Sum of fipronil, the sulphenyl metabolite (5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl) sulphenyl]-1H-pyrazole-3-carbonitrile), the sulphonyl metabolite (5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl)sulphonyl]-1H-pyrazole-3-carbonitrile), and the trifluoromethyl metabolite (5-amino-4-trifluoromethyl-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1H-pyrazole-3-carbonitrile)		Sweet potato	*0.01
Asparagus	0.2	Turnip, garden	0.1
Assorted tropical and sub-tropical fruit	T*0.01	Wine grapes	*0.01
– inedible peel [except banana; custard apple]		Flamprop-methyl	
Banana	0.01	Flamprop-methyl	
Bergamot	T0.1	Edible offal (mammalian)	*0.01
Brassica (cole or cabbage) vegetables,	T0.05	Lupin (dry)	0.05
		Meat (mammalian)	*0.01
		Milks	*0.01
		Safflower seed	*0.05
		Triticale	0.05
		Wheat	0.05
		Flamprop-M-methyl	
		see Flamprop-methyl	
		Flavophospholipol	
		Flavophospholipol	
		Cattle fat	*0.01

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Cattle kidney	*0.01	Eggs	*0.05
Cattle liver	*0.01	Fruiting vegetables, cucurbits	0.1
Cattle meat	*0.01	Galangal, rhizomes	0.05
Cattle milk	T*0.01	Garlic	0.05
Eggs	*0.02	Ginger, root	0.05
Flonicamid		Herbs	T2
Flonicamid [N -(cyanomethyl)-4-(trifluoromethyl)-3-pyridinecarboxamide] and its metabolites TFNA [4-trifluoromethylnicotinic acid], TFNA-AM [4-trifluoromethylnicotinamide] TFNG [N -(4-trifluoromethylnicotinoyl)glycine]		Hops, dry	0.05
Cotton seed	1	Leafy vegetables [except lettuce, head]	T2
Edible offal (mammalian)	*0.02	Leek	T1
Eggs	*0.02	Legume vegetables	0.1
Fruiting vegetables, cucurbits	0.7	Lettuce, head	0.05
Meat (mammalian)	*0.02	Lotus root	T3
Milks	*0.02	Lupin (dry)	0.1
Pome fruits	0.7	Meat (mammalian)	*0.05
Potato	0.2	Milks	0.1
Poultry, edible offal of	*0.02	Oilseed	0.5
Poultry meat	*0.02	Onion, bulb	0.05
Stone fruits	0.6	Onion, Chinese	0.05
Florasulam		Onion, Welsh	0.05
Florasulam		Peppers, Sweet	*0.02
Cereal grains	*0.01	Pome fruits	*0.01
Edible offal (mammalian)	*0.01	Potato	0.05
Eggs	*0.01	Poultry, edible offal of	*0.05
Meat (mammalian)	*0.01	Poultry meat	*0.05
Milks	*0.01	Pulses	0.5
Poultry, edible offal of	*0.01	Root and tuber vegetables [except potato; sweet potato; taro; yam bean; yams]	T1
Poultry meat	*0.01	Shallot	0.05
Florfenicol		Spring Onion	0.05
Sum of florfenicol and its metabolites florfenicol alcohol, florfenicol oxamic acid, monochloroflorfenicol and florfenicol amine expressed as florfenicol amine		Stone fruits	0.05
Cattle kidney	0.5	Sugar cane	T*0.1
Cattle liver	3	Sweet potato	T0.3
Cattle meat	0.3	Taro	T3
Fish	T0.5	Tea, green, black	T50
Pig fat/skin	1	Tomato	0.1
Pig kidney	1	Turmeric, root	0.05
Pig liver	3	Water chestnut	T3
Pig meat	0.5	Yam bean	T3
Fluazifop-p-butyl		Yams	T0.3
Sum of fluazifop-butyl, fluazifop and their conjugates, expressed as fluazifop		Fluazinam	
Assorted tropical and sub-tropical fruits — inedible peel [except avocado and banana]	0.05	Fluazinam	
Avocado	*0.02	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	*0.01
Banana	*0.02	Pome fruits	*0.01
Berries and other small fruits	0.2	Potato	*0.01
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	1	Wine grapes	*0.05
Celery	*0.02	Fluazuron	
Chia	T2	Fluazuron	
Citrus fruits	*0.02	Cattle, edible offal of	0.5
Coriander (leaves, stem, roots)	T2	Cattle meat (in the fat)	7
Date	T0.2	Flubendiamide	
Edible offal (mammalian)	*0.05	Commodities of plant origin: Flubendiamide Commodities of animal origin: Sum of flubendiamide and 3-iodo-N-(2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl)phthalimide, expressed as flubendiamide	
Egg plant	T0.7	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	5
		Chia	1
		Common bean (pods and/or immature	T2

Maximum Residue Limits (mg/kg)

Standard 1.4.2

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Flunixin Flunixin			Fluroxypyr Fluroxypyr		
Cattle kidney	0.02		Cereal grains		0.2
Cattle liver	0.02		Edible offal (mammalian) [except kidney]		0.1
Cattle meat (in the fat)	0.02		Eggs		*0.01
Fluometuron Sum of fluometuron and 3-trifluoromethylaniline, expressed as fluometuron			Kidney (mammalian)		1
Cereal grains	*0.1		Meat (mammalian) (in the fat)		0.1
Citrus fruits	0.5		Milks		0.1
Cotton seed	*0.1		Poultry, edible offal of		*0.05
Pineapple	*0.1		Poultry meat		*0.05
Fluopicolide Fluopicolide			Sugar cane (in the juice)		0.2
Grapes	2		Sweet corn (corn-on-the-cob)		0.2
Fluopyram <i>Commodities of plant origin:</i> Fluopyram <i>Commodities of animal origin for enforcement:</i> Sum of fluopyram and 2-(trifluoromethyl)-benzamide, expressed as fluopyram <i>Commodities of animal origin for dietary exposure assessment:</i> Sum of fluopyram, 2-(trifluoromethyl) benzamide and the combined residues of <i>N</i> -{(E)-2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethenyl}-2-(trifluoromethyl) benzamide and <i>N</i> -{(Z)-2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethenyl}-2-(trifluoromethyl) benzamide, all expressed as fluopyram			Flusilazole Flusilazole		
Almonds	T0.5		Grapes		0.5
Banana	T0.1		Pome fruits		0.2
Cherries	T5		Sugar cane		*0.02
Dried grapes (currants, raisins and sultanas)	T15		Flutolanil <i>Commodities of plant origin:</i> Flutolanil <i>Commodities of animal origin:</i> Flutolanil and metabolites hydrolysed to 2-trifluoromethyl-benzoic acid and expressed as flutolanil		
Edible offal (mammalian)	T0.7		Edible offal (mammalian)		*0.05
Meat (mammalian)	T0.05		Eggs		*0.05
Milks	T0.2		Meat (mammalian) (in the fat)		*0.05
Pome fruits	T0.5		Milks		*0.05
Stone fruits [except cherries]	T2		Potato		0.05
Table-grapes	T2		Poultry, edible offal of		*0.05
Fluoxastrobin Sum of fluoxastrobin and its Z isomer			Poultry meat (in the fat)		*0.05
Cranberry	1.9		Flutriafol Flutriafol		
Flupropanate Flupropanate			Barley		0.2
Edible offal (mammalian)	*0.1		Cereal grains [except as otherwise listed under this chemical]		*0.02
Meat (mammalian) (in the fat)	*0.1		Edible offal (mammalian)		0.5
Milks	0.1		Eggs		*0.05
Fluquinconazole Fluquinconazole			Garden pea (young pods)		*0.01
Barley	*0.02		Meat (mammalian)		*0.05
Edible offal (mammalian)	0.2		Milks		*0.05
Eggs	*0.02		Poultry, edible offal of		*0.05
Meat (mammalian) (in the fat)	0.5		Poultry meat		*0.05
Milks	*0.02		Rape seed (canola)		*0.02
Pome fruits	0.3		Sugar cane		*0.01
Poultry, edible offal of	*0.02		Fluvalinate Fluvalinate, sum of isomers		
Poultry meat (in the fat)	*0.02		Apple		0.1
Rape seed (canola)	*0.01		Asparagus		0.2
Wheat	*0.02		Cauliflower		0.5
			Cotton seed		0.1
			Honey		T*0.01
			Stone fruits		0.05
			Table grapes		0.05
			Tomato		0.5
			Fluxapyroxad <i>Commodities of plant origin:</i> Fluxapyroxad <i>Commodities of animal origin for enforcement:</i> Fluxapyroxad		
			All other foods		0.1

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Barley	0.2	Eggs	*0.05
Barley bran, unprocessed	0.5	Hops, dry	T1
Edible offal (mammalian)	0.03	Lemon myrtle	T20
Eggs	0.005	Maize	0.2
Meat (mammalian) (in the fat)	0.05	Meat (mammalian)	0.1
Milk fats	0.02	Milks	*0.05
Milks	0.005	Native foods [except lemon myrtle]	T0.1
Poultry, edible offal of	*0.01	Oilseeds [except cotton seed; rape seed (canola)]	*0.1
Poultry meat (in the fat)	*0.01	Olives	*0.1
Fluxapyroxad Fluxapyroxad		Peppers, Sweet (capsicum)	*0.05
Plums (including prunes)	3	Podded pea (young pods) (snow and sugar snap)	T1
Pome fruits	0.8	Pome fruits	*0.1
Pulses [except soya bean (dry)]	0.4	Poultry, edible offal of	*0.1
Soya bean (dry)	0.3	Poultry meat	*0.05
Soya bean (immature seeds)	0.15	Pulses [except soya bean (dry)]	*0.1
Stone fruits [except plums (including prunes)]	2	Rape seed (canola)	5
Forchlorfenuron Forchlorfenuron		Saffron	T*0.05
Blueberries	T*0.01	Soya bean (dry)	2
Grapes	*0.01	Stone fruits	*0.05
Kiwifruit	T*0.01	Sugar cane	T*0.2
Mango	T*0.01	Tomato	*0.05
Plums (including prunes)	T*0.01	Tea, green, black	T20
Prunes	T*0.01	Tree nuts	0.1
Fosetyl Fosetyl		Glyphosate Sum of glyphosate and Aminomethylphosphonic acid (AMPA) metabolite, expressed as glyphosate	
Apple	1	Adzuki bean (dry)	10
Avocado	5	Avocado	*0.05
Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	T0.1	Babaco	*0.05
Durian	T5	Banana	0.2
Fruiting vegetables, other than cucurbits	T0.02	Barley	10
Leafy vegetables [except rucola (rocket); spinach]	T0.2	Berries and other small fruits	*0.05
Peach	1	Bulb vegetables	*0.1
Pineapple	5	Cereal grains [except barley; maize; sorghum; wheat]	T*0.1
Rucola (rocket)	T0.7	Citrus fruits	0.5
Spinach	T0.7	Coffee beans	T0.2
Stone fruits [except cherries; peach]	T1	Cotton seed	15
Furathiocarb see Carbofuran Residues arising from the use of furathiocarb are covered by MRLs for carbofuran		Cotton seed oil, crude	*0.1
Glufosinate and Glufosinate-ammonium Sum of glufosinate-ammonium, N-acetyl glufosinate and 3-[hydroxy(methyl)-phosphinoyl] propionic acid, expressed as glufosinate (free acid)		Cowpea (dry)	10
Assorted tropical and sub-tropical fruits – inedible peel	0.2	Custard apple	*0.05
Berries and other small fruits	0.1	Date	T2
Cereal grains	*0.1	Edible offal (mammalian)	2
Citrus fruits	0.1	Eggs	*0.05
Coffee beans	T*0.05	Fig	*0.05
Cotton seed	3	Fruiting vegetables, cucurbits	*0.1
Date	T0.1	Fruiting vegetables, other than cucurbits	*0.1
Edible offal (mammalian)	5	Guar bean (dry)	10
		Guava	*0.05
		Hops, dry	*0.1
		Kiwifruit	*0.05
		Leafy vegetables	*0.1
		Legume vegetables	*0.1
		Lemon myrtle	T20
		Linseed	T5
		Litchi	0.2
		Maize	5
		Mango	*0.05
		Meat (mammalian)	*0.1
		Milks	*0.1
		Monstero	*0.05
		Mung bean (dry)	10

Maximum Residue Limits (mg/kg)

Federal Register of Legislative Instruments F2015C00185

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Hydrogen phosphide see Phosphine		Pulses	*0.1
Imazalil Imazalil		Imidacloprid Sum of imidacloprid and metabolites containing the 6-chloropyridinylmethylene moiety, expressed as imidacloprid	
Chicken, edible offal of	*0.01	Apple	0.3
Chicken meat	*0.01	Assorted tropical and sub-tropical fruits – inedible peel [except banana]	T1
Citrus fruits	10	Banana	0.5
Eggs	*0.01	Beetroot	T0.05
Melons, except watermelon	10	Bergamot	T5
Mushrooms	T1	Berries and other small fruits [except blueberries; cranberry; grapes; strawberry]	5
Pome fruits	5	Blueberries	T0.1
Potato	5	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	0.5
Imazamox Imazamox		Broad bean (dry)	*0.05
Adzuki bean (dry)	T*0.05	Burdock, greater	T0.05
Barley	*0.05	Burnet, Salad	T5
Broad bean (dry) (fava beans)	T*0.05	Cereal grains [except maize; popcorn; sorghum]	*0.05
Edible offal (mammalian)	*0.05	Celery	0.3
Field pea (dry)	*0.05	Citrus fruits	2
Meat (mammalian)	*0.05	Common bean (dry) (navy bean)	T1
Milks	*0.05	Common bean (pods and/or immature seeds)	T1
Peanut	*0.05	Coriander (leaves, stem, roots)	T5
Poppy seed	T*0.05	Coriander, seed	T5
Rape seed (canola)	*0.05	Cotton seed	*0.02
Soya bean (dry)	*0.05	Date	T1
Wheat	*0.05	Dill, seed	T5
Imazapic Sum of imazapic and its hydroxymethyl derivative		Edible offal (mammalian)	0.2
Edible offal (mammalian)	*0.05	Eggs	*0.02
Eggs	*0.01	Fennel, bulb	T0.1
Meat (mammalian) (in the fat)	*0.05	Fennel, seed	T5
Milks	*0.01	Field pea (dry)	*0.05
Peanut	*0.1	Fruiting vegetables, cucurbits	0.2
Poultry, edible offal of	*0.01	Fruiting vegetables, other than cucurbits [except sweet corn, (corn-on-the-cob)]	0.5
Poultry meat	*0.01	Galangal, Greater	T0.05
Rape seed (canola)	*0.05	Garlic	T0.5
Sugar cane	*0.05	Ginger, Japanese	T5
Wheat	*0.05	Ginger, root	T0.3
Imazapyr Imazapyr		Grapes	T0.1
Barley	*0.05	Hazelnuts	T*0.01
Edible offal (mammalian)	*0.05	Herbs	T5
Meat (mammalian) (in the fat)	*0.05	Hops, dry	T10
Maize	*0.05	Kaffir lime leaves	T5
Milks	*0.01	Leafy vegetables [except lettuce, head]	20
Poppy seed	T*0.05	Lemon balm	T5
Rape seed (canola)	*0.05	Lemon grass	T5
Wheat	*0.05	Lemon verbena (fresh weight)	T5
Imazethapyr Imazethapyr		Lentil (dry)	0.2
Edible offal (mammalian)	*0.1	Lettuce, head	5
Eggs	*0.1	Lupin (dry)	0.2
Legume vegetables	*0.1	Maize	0.05
Maize	*0.05	Meat (mammalian)	0.05
Meat (mammalian)	*0.1	Milks	0.05
Milks	*0.1	Peanut	T0.5
Peanut	*0.1	Persimmon, Japanese	T1
Poultry, edible offal of	*0.1	Popcorn	0.05
Poultry meat	*0.1	Potato	0.3

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Poultry, edible offal of	*0.02	Safflower seed	T0.5
Poultry meat	*0.02	Stone fruits	2
Radish, Japanese	T0.05	Sunflower seed	T1
Rape seed (canola)	*0.05	Tomato	T0.5
Rhubarb	T0.2		
Rose and dianthus (edible flowers)	T5		
Sorghum	*0.02		
Stone fruits	0.5		
Strawberry	0.5		
Sugar cane	*0.05		
Sunflower seed	*0.02		
Sweet corn (corn-on-the-cob)	*0.05		
Sweet potato	0.3		
Taro	T0.05		
Teas (tea and herb teas)	T10		
Tree tomato	T2		
Turmeric, root (fresh)	T0.05		
Yam bean	T0.05		
Yams	T0.05		
Imidocarb (dipropionate salt) Imidocarb		Inorganic bromide Bromide ion	
Cattle, edible offal of	5	Avocado	75
Cattle meat	1	Cereal grains	50
Cattle milk	0.2	Citrus fruits	30
		Dates, dried	100
		Dried fruits [except as otherwise listed under this chemical]	30
		Dried grapes	100
		Dried herbs	400
		Dried peach	50
		Figs, dried	250
		Fruit [except as otherwise listed under this chemical]	20
		Peppers, Sweet	50
		Prunes	20
		Spices	400
		Strawberry	30
		Vegetables [except as otherwise listed under this chemical]	20
Indoxacarb Sum of indoxacarb and its <i>R</i> -isomer		Iodosulfuron methyl Iodosulfuron methyl	
Asparagus	T1	Barley	*0.01
Berries and other small fruits [except grapes]	T1	Edible offal (mammalian)	*0.01
Brassica (cole or cabbage) vegetables, Head cabbages and Flowerhead brassicas	2	Eggs	*0.01
Celery	T5	Meat (mammalian) (in the fat)	*0.01
Chervil	T10	Milks	*0.01
Chia	T*0.05	Poultry, edible offal of	*0.01
Coriander (leaves, stem, roots)	T20	Poultry meat (in the fat)	*0.01
Cotton seed	1	Wheat	*0.01
Dried grapes	2		
Edible offal (mammalian) [except kidney]	*0.01		
Egg plant	0.5		
Eggs	*0.01		
Grapes	0.5		
Herbs	T20		
Kidney (mammalian)	0.2		
Leafy vegetables [except chervil; lettuce, head; mizuna; rucola]	5		
Lemon balm	T10		
Lettuce, head	3		
Linseed	T0.5		
Meat (mammalian) (in the fat)	1		
Mexican tarragon	T20		
Milk fats	1		
Milks	0.01		
Mizuna	T10		
Olives	T0.2		
Peanut	T0.02		
Peppers, Sweet	0.5		
Pome fruits	2		
Poultry (edible offal of)	*0.01		
Poultry meat (in the fat)	*0.01		
Pulses	0.2		
Rape seed (canola)	T*0.05		
Rucola (rocket)	T20		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Brassica leafy vegetables	15		
Broad bean (green pods and immature seeds)	0.2		
Broccoli	T*0.05		
Brussels sprouts	0.5		
Cabbages, head	T*0.05		
Carrot	T0.5		
Cauliflower	T*0.05		
Celeriac	T0.7		
Celery	2		
Chard (silver beet)	T15		
Edible offal (mammalian)	*0.1		
Egg plant	T1		
Garlic	T10		
Grapes	20		
Kiwifruit	10		
Lettuce, head	5		
Lettuce, leaf	5		
Lupin (dry)	*0.1		
Macadamia nuts	*0.01		
Mandarins	T5		
Meat (mammalian)	*0.1		
Milks	*0.1		
Onion, bulb	T0.7		
Passionfruit	10		
Peanut	0.05		
Peanut oil, crude	0.05		
Peppers	T3		
Pistachio nut	T*0.05		
Pome fruits	3		
Potato	*0.05		
Rape seed (canola)	0.5		
Soya bean (dry)	0.05		
Spinach	T5		
Stone fruits	10		
Tangelo, large-sized cultivars	T5		
Tomato	2		
Isoeugenol			
Isoeugenol, sum of cis- and trans- isomers			
Diadromous fish (whole commodity)	100		
Freshwater fish (whole commodity)	100		
Marine fish (whole commodity)	100		
Isoxaben			
Isoxaben			
Assorted tropical and sub-tropical fruits – edible peel	*0.01		
Assorted tropical and sub-tropical fruits – inedible peel	*0.01		
Barley	*0.01		
Citrus fruits	*0.01		
Edible offal (mammalian)	*0.01		
Eggs	*0.01		
Grapes	*0.01		
Hops, dry	*0.1		
Meat (mammalian)	*0.01		
Milks	*0.01		
Pome fruits	*0.01		
Poultry, edible offal of	*0.01		
Poultry meat	*0.01		
Stone fruits	*0.01		
Tree nuts	*0.01		
Triticale	*0.01		
Wheat	*0.01		
Isoxaflutole			
The sum of isoxaflutole and 2-cyclopropylcarbonyl-3-(2-methylsulfonyl-4-trifluoromethylphenyl)-3-oxopropanenitrile, expressed as isoxaflutole			
Cereal grains	*0.02		
Chick-pea (dry)	*0.02		
Edible offal (mammalian)	0.1		
Eggs	*0.05		
Meat (mammalian)	*0.05		
Milks	*0.05		
Poppy seed	*0.02		
Poultry, edible offal of	*0.05		
Poultry meat	*0.05		
Sugar cane	*0.01		
Ivermectin			
H ₂ B _{1a}			
Cattle kidney	*0.01		
Cattle liver	0.1		
Cattle meat (in the fat)	0.04		
Cattle milk	0.05		
Deer kidney	*0.01		
Deer liver	*0.01		
Deer meat (in the fat)	*0.01		
Horse, edible offal of	*0.01		
Horse meat	*0.01		
Pig kidney	*0.01		
Pig liver	*0.01		
Pig meat (in the fat)	0.02		
Sheep kidney	*0.01		
Sheep liver	0.015		
Sheep meat (in the fat)	0.02		
Ketoprofen			
Ketoprofen			
Cattle, edible offal of	*0.05		
Cattle meat	*0.05		
Cattle milk	*0.05		
Kitasamycin			
Inhibitory substance, identified as kitasamycin			
Eggs	*0.2		
Pig, edible offal of	*0.2		
Pig meat	*0.2		
Kresoxim-methyl			
<i>Commodities of plant origin:</i> Kresoxim-methyl			
<i>Commodities of animal origin:</i> Sum of a-(p-hydroxy-o-tolyloxy)-o-tolyl (methoxyimino) acetic acid and (E)-methoxyimino[a-(o-tolyloxy)-o-tolyl]acetic acid, expressed as kresoxim-methyl			
Edible offal (mammalian)	*0.01		
Fruiting vegetables, cucurbits	0.05		
Grapes	1		
Meat (mammalian)	*0.01		
Milks	*0.001		
Pome fruits	0.1		
Lambda-cyhalothrin			
see Cyhalothrin			
Lasalocid			
Lasalocid			
Cattle milk	*0.01		
Edible offal (mammalian)	0.7		

SCHEDULE 1

Maximum Residue Limits (mg/kg)

Eggs	*0.05	Milks	T0.2
Meat (mammalian)	*0.05	Poultry, edible offal of	T*0.01
Poultry, edible offal of	0.4	Poultry meat (in the fat)	T1
Poultry meat	*0.1		
Poultry skin/fat	1		
Levamisole			
Levamisole			
Edible offal (mammalian)	1		
Eggs	1		
Goat milk	0.1		
Meat (mammalian)	0.1		
Milks [except goat milk]	0.3		
Poultry, edible offal of	0.1		
Poultry meat	0.1		
Lincomycin			
Inhibitory substance, identified as lincomycin			
Cattle milk	*0.02		
Edible offal (mammalian) [except sheep, edible offal of]	0.2		
Eggs	0.2		
Goat milk	*0.1		
Meat (mammalian) [except sheep meat]	0.2		
Poultry, edible offal of	0.1		
Poultry meat	0.1		
Lindane			
Lindane			
Pineapple	0.5		
Linuron			
Sum of linuron plus 3,4-dichloroaniline, expressed as linuron			
Celeriac	T0.5		
Celery	*0.05		
Cereal grains	*0.05		
Chervil	T1		
Coriander (leaves, stem, roots)	T1		
Coriander, seed	0.2		
Edible offal (mammalian)	1		
Eggs	*0.05		
Herbs	T1		
Leek	*0.02		
Lemon grass	T1		
Lemon verbena (dry leaves)	T1		
Meat (mammalian)	*0.05		
Milks	*0.05		
Mizuna	T1		
Parsnip	T0.05		
Poultry, edible offal of	*0.05		
Poultry meat	*0.05		
Rucola (rocket)	T1		
Turmeric root	T*0.05		
Vegetables [except celeriac; celery; leek; parsnip]	*0.05		
Lufenuron			
Lufenuron			
Cotton seed	T0.2		
Cotton seed oil, crude	T0.5		
Edible offal (mammalian)	T*0.01		
Eggs	T0.05		
Meat (mammalian) (in the fat)	T1		