Schedule 1—Specified permissible ingredients and requirements applying to these ingredients when contained in a medicine

Note: See sections 5 and 6.

Column 1	Column 2	Column 3	Column 4
Item	Ingredient Name	Purpose	Specific requirements
5067	UBIDECARENONE	Α, Ε	When used as an excipient, the route of administration must be topical and the concentration in the medicine must not be more than 0.05%.
			Not to be included in medicines intended for use in the eye.
			When for internal use, the maximum recommended daily dose must not provide more than 300 milligrams of ubidecarenone.
			When for internal use in combination with Ubiquinol-10, the maximum recommended daily dose must not provide more than 300 milligrams of ubiquinol-10 and ubidecarenone combined.
			When for internal use, the following warning statement is required on the medicine label:
			- (WARF) 'Do not take while on warfari therapy without medical advice'.
5068	UBIQUINOL-10	Α, Ε	When used as an excipient, the route of administration must be topical and the concentration in the medicine must be no more than 0.05%.
			Not to be included in medicines intended for use in the eye.
			When for internal use, the maximum recommended daily dose must provide no more than 300 milligrams of ubiquinol-10.
			When used in combination with ubidecarenone, the maximum

Volume 6

recommended daily dose must provide no more than 300 mg of ubiquinol-10 and ubidecarenone combined.

The medicine requires the following warning statement on the medicine label:

- (WARF) 'Do not take while on warfarin therapy without medical advice.'

5069	ULEX EUROPAEUS	A, H	
5070	ULMUS AMERICANA	A, H	
5071	ULMUS CAMPESTRIS	A, H	
5072	ULMUS GLABRA	A, H	
5073	ULMUS MINOR	A, H	
5074	ULMUS PARVIFOLIA	A, H	
5075	ULMUS PUMILA	A, H	
5076	ULMUS RUBRA	A, H	
5077	ULTRALIDE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
5078	ULTRAMARINE BLUE	E	Permitted for use only as a colour for topical use.
5079	ULVA LACTUCA	А, Н	Iodine is a mandatory component of Ulva lactuca.
			Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 0.1%.
5080	UMBELLULARIA CALIFORNICA	A, H	
5081	UNCARIA GAMBIR	A, H	
5082	UNCARIA RHYNCOPHYLLA	A, H	
5083	UNCARIA SINENSIS	A, H	
5084	UNCARIA TOMENTOSA	A, H	
5085	UNDARIA PINNATIFIDA	А, Н	Whole dried Undaria pinnatifida must not contain the holdfast.

Vol	ume	6
-----	-----	---

			Only for use in oral medicines.
5086	UNDECANAL	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
5087	UNDECANOIC ACID	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used as a flavour the total flavour concentration in a medicine must be no more than 5%.
5088	UNDECENOIC ACID	Е	
5089	UNDECYL ALCOHOL	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
5090	UNDECYLCRYLENE DIMETICONE	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 10%.
5091	UNDECYLENAMIDE DEA	E	
5092	UNDECYLENOYL PEG-5 PARABEN	Е	Only for use in topical medicines for dermal application.
5093	URANIUM NITRATE	Н	Only for use as an active homoeopathic ingredient.

Volume 6

5094	UREA	А, Е, Н	Only for use in topical medicines for dermal application.
			The concentration in the medicine must be no more than 10% (w/w).
5095	URTICA DIOICA	A, E, H	
5096	URTICA URENS	A, H	
5097	USNEA BARBATA	A, H	
5098	UVA URSI LEAF DRY	A, H	
5099	UVA URSI LEAF POWDER	А, Е, Н	
5100	VA/BUTYL MALEATE/ISOBORNYL ACRYLATE COPOLYMER	Е	Vinyl acetate is a mandatory component of VA/butyl maleate/isobornyl acrylate copolymer.
			The concentration of vinyl acetate in the medicine must be no more than 0.01% or 100 ppm.
			Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 5%.
5101	VACCARIA SEGATALIS	A, H	
5102	VACCINIUM BRACTEATUM	A, H	
5103	VACCINIUM CORYMBOSUM	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5104	VACCINIUM MACROCARPON	A, E, H	
5105	VACCINIUM MYRTILLOIDES	A, H	
5106	VACCINIUM MYRTILLUS	А, Е, Н	
5107	VACCINIUM OXYCOCCUS	A, H	
5108	VACCINIUM VITIS-IDAEA	А, Н	Beta-arbutin is a mandatory component of Vaccinium vitis-idaea.
			When for oral use, the maximum recommended daily dose must not

			provide more than 500 mg of beta- arbutin.
			When for dermal application exclusively to the face:
			a) the concentration of beta-arbutin in the medicine must not be more than 7%;
			b) hydroquinone is a mandatory component; and
			c) the concentration of hydroquinone must not be more than 10 mg/kg or 10 mg/L or 0.001%.
			When for use other than oral or dermal application exclusively to the face, the concentration of beta-arbutin in the medicine must not be more than 10 mg/kg or 10 mg/L or 0.001%.
5109	VALENCENE	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5110	VALERALDEHYDE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5111	VALERIAN DRY	А, Н	The following warning statement is required on the medicine label when the medicine is for oral use:
			(VALER) 'In rare cases, valerian may harm the liver. Stop use and see a doctor if you have yellowing skin/eyes or unusual: fatigue, nausea, appetite loss, abdominal pain, dark urine or itching.'
5112	VALERIAN OIL	Е	Permitted for use only in combination with other permitted ingredients as a flavour.

Volume 6

			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5113	VALERIAN POWDER	А, Н	The following warning statement is required on the medicine label when the medicine is for oral use:
			(VALER) 'In rare cases, valerian may harm the liver. Stop use and see a doctor if you have yellowing skin/eyes or unusual: fatigue, nausea, appetite loss, abdominal pain, dark urine or itching.'
5114	VALERIANA EDULIS	A, H	
5115	VALERIANA OFFICINALIS	А, Н	The following warning statement is required on the medicine label when the medicine is for oral use:
			(VALER) 'In rare cases, valerian may harm the liver. Stop use and see a doctor if you have yellowing skin/eyes or unusual: fatigue, nausea, appetite loss, abdominal pain, dark urine or itching.'
5116	VALERIANA SORBIFOLIA	A, H	
5117	VALERIC ACID	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5118	VALINE	A, E	
5119	VANADIUM	Н	
5120	VANILLA	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.

Vol	lume	6
101		v

5121	VANILLA DRY	А, Е, Н	
5122	VANILLA EXTRACT	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5123	VANILLA OLEORESIN	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5124	VANILLA PLANIFOLIA	A, E, H	
5125	VANILLA POWDER	A, E, H	
5126	VANILLA TAHITENSIS	A, H	
5127	VANILLIC ACID	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5128	VANILLIN	Е	
5129	VANILLIN ACETATE	Е	Vanillin acetate must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation.
			The total concentration of flavour proprietary excipient formulations containing vanillin acetate must not be more than 5% of the total medicine.
			The maximum recommended daily dose of the medicine must not provide more than 1.8 micrograms of vanillin acetate.
5130	VANILLIN ISOBUTYRATE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.

Volume 6

			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
5131	VANILLYL ALCOHOL	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5132	VAT RED 1	E	Permitted for use only as a colour for topical use.
5133	VAT RED 1 ALUMINIUM LAKE	E	Permitted for use only as a colour for topical use.
5134	VAT RED 5	E	Permitted for use only as a colour for topical use.
5135	VEGETABLE OIL	Е	
5136	VEGETABLE OIL PHYTOSTEROL	А	Only for use in oral medicines.
	ESTERS		The medicine requires the following warning statement on the medicine label
			- (PREGNT) 'Not recommended for use by pregnant and lactating women (or words to that effect).'
5137	VEIN	Н	Only for use as an active homoeopathic ingredient.
5138	VERATRALDEHYDE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Vo	lume	6
V U.	luine	v

			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
5139	VERATROL	E	Permitted for use only in combination with other permitted ingredients as part of a fragrance proprietary excipient formulation.
			The total fragrance proprietary excipient formulation in a medicine must be no more than 1%.
5140	VERATRUM ALBUM	A, H	Solanidine is a mandatory component of Veratrum album.
			The concentration of equivalent dry Veratrum album in the medicine must be no more than 10mg/Kg or 10mg/L or 0.001%.
5141	VERBASCUM DENSIFLORUM	A, H	
5142	VERBASCUM THAPSUS	A, H	
5143	VERBENA OFFICINALIS	A, H	
5144	VERBENA OIL	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5145	VERONICA CHAMAEDRYS	A, H	
5146	VERONICA OFFICINALIS	A, H	
5147	VERONICASTRUM VIRGINICUM	A, E, H	
5148	VERTONAL	E	Permitted for use only in combination with other permitted ingredients as part of a fragrance proprietary excipient formulation.
			When included in a medicine for use on the lips the concentration of vertonal must be no more than 0.2%.
			The total fragrance proprietary excipient formulation in a medicine must be no more than 1%.

Volume 6

5149	VETIVER OIL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
5150	VETIVERYL ACETATE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
5151	VIBURNUM OPULUS	A, E, H	
5152	VIBURNUM PRUNIFOLIUM	A, E, H	
5153	VICIA FABA	А, Н	Levodopa is a mandatory component of Vicia faba.
			The concentration of levodopa in the medicine must not be more than 10 mg/kg or 10 mg/L or 0.001%.
5154	VIGNA ANGULARIS VAR. ANGULARIS	A, H	
5155	VIGNA RADIATA	A, H	
5156	VIGNA UMBELLATA	A, H	
5157	VINCA MAJOR	A, H	Vincamine is a mandatory component of Vinca major.
			The concentration of vincamine in the medicine must be no more than 10mg/kg or 10 mg/L or 0.001%.
5158	VINCA MINOR	A, H	Vincamine and vincristine are mandatory components of Vinca minor.
			The concentration of vincamine in the medicine must be no more than 10mg/kg or 10 mg/L or 0.001%.

_

			The concentration of Vincristine in the medicine must be no more than 10mg/kg or 10mg/L or 0.001%
5159	VINCETOXICUM OFFICINALE	A, H	
5160	VINEGAR	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5161	VIOLA ODORATA	A, E, H	
5162	VIOLA TRICOLOR	А, Н	
5163	VIOLA YEDOENSIS	A, H	
5164	VIOLET LEAF ABSOLUTE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
5165	VIPER	Н	Only for use as an active homoeopathic ingredient.
5166	VISCUM ALBUM	A, E, H	
5167	VISCUM COLORATUM	А, Н	
5168	VISCUM FLAVESCENS	А, Н	
5169	VITELLARIA PARADOXA	А, Е, Н	
5170	VITEX AGNUS-CASTUS	А, Е, Н	When the ingredient is in a medicine tha is for internal use, the following warning statement is required on the label:
			 - (VAC) 'Vitex agnus-castus may affect hormones and medicines such as oral contraceptives. Consult your health professional before use' (or words to that effect).

Volume 6

5171	VITEX NEGUNDO	A, H	
5172	VITEX ROTUNDIFOLIA	A, H	
5173	VITEX TRIFOLIA	A, H	
5174	VITIS VINIFERA	А, Е, Н	
5175	VITREOSCILLA CONCENTRATE	Е	Only for use in topical medicines for dermal application.
			The concentration in the medicine must be no more than 0.1%.
5176	VP/ACRYLATES/LAURYL METHACRYLATE COPOLYMER	Е	Only for use in topical medicines for dermal application and not to be included in topical medicines intended for use in the eye.
			The concentration in the medicine must not be more than 2.00%.
5177	WAHLENBERGIA GRACILIS	A, H	
5178	WALNUT	Е	
5179	WALNUT OIL	Е	
5180	WATER MELON	Е	
5181	WHEAT	Е	Gluten is a mandatory component of Wheat when the route of administration is other than topical and mucosal.
5182	WHEAT BRAN	Е	Gluten is a mandatory component of Wheat bran when the route of administration is other than topical and mucosal.
5183	WHEAT DEXTRIN	Α, Ε	Gluten is a mandatory component of wheat dextrin.
			Only for use when the dosage form is capsule, tablet or pill.
5184	WHEAT GERM	E	Gluten is a mandatory component of Wheat germ when the route of administration is other than topical and mucosal.
5185	WHEAT GERM GLYCERIDES	Е	Gluten is a mandatory component of wheat germ glycerides when the route of

			administration is other than topical and mucosal.
5186	WHEAT LEAF	E	
5187	WHEAT STARCH	Е	When the route of administration is other than topical or mucosal, gluten is a mandatory component of wheat starch.
5188	WHEATGERM OIL	A, E, H	
5189	WHEY POWDER	Е	Lactose is a mandatory component of Whey powder when the route of administration is oral.
5190	WHEY PROTEIN	Е	Lactose is a mandatory component of Whey protein when the route of administration is oral.
5191	WHEY PROTEIN CONCENTRATE	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5192	WHITE BEESWAX	E	
5193	WHITE HOREHOUND HERB DRY	A, H	
5194	WHITE HOREHOUND HERB POWDER	А, Н	
5195	WHITE SOFT PARAFFIN	A, E	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing from time to time.
5196	WHOLE DRY MILK	E	
5197	WIKSTROEMIA VIRIDIFLORA	А, Н	
5198	WILD CARROT HERB DRY	А, Е, Н	
5199	WILD CARROT HERB POWDER	A, H	
5200	WILD CHERRY BARK DRY	A, H	

Volume	6
--------	---

5201	WILD CHERRY BARK POWDER	А, Н	
5202	WILD LETTUCE LEAF DRY	A, H	
5203	WILD LETTUCE LEAF POWDER	A, H	
5204	WINTERGREEN OIL	А, Е, Н	Methyl salicylate is a mandatory component of wintergreen oil.
			Not to be included in medicines for use in the eye or on damaged skin.
			When used internally, the concentration of methyl salicylate in the medicine must not be more than 0.001%.
			When the concentration of methyl salicylate in a liquid preparation is more than 5% and the dosage form is other than spray, the medicine requires child resistant packaging.
			When the concentration of methyl salicylate in a liquid preparation is more than 5%, and the dosage form is spray, the medicine does not require child resistant packaging if:
			 the delivery device is engaged into the container in such a way that prevents it from being readily removed;
			 direct suction through the delivery device results in delivery of no more that one dosage unit; and
			 actuation of the spray device is ergonomically difficult for young children to accomplish.
			The following warning statement is required on the medicine label:
			- (METSAL) 'Contains methyl salicylat (or words to that effect).
			When for use in topical medicines for dermal application:
			i) the concentration of methyl salicylatein the medicine must not be more than25%;
			ii) the following warning statements are required on the medicine label:
			- (PREGNT2) 'Do not use if pregnant or likely to become pregnant' (or words to that effect);

			Volume 6
			- (CHILD4) 'Do not use [this product/insert name of product] in children 6 years of age or less';
			- (SENS) 'Application to skin may increase sensitivity to sunlight' (or words to that effect);
			- (AVOID) 'Avoid prolonged exposure in the sun' (or words to that effect);
			iii) if the concentration of methyl salicylate in the medicine is greater than 1%, the following warning statement is required on the medicine label:
			- (IRRIT) 'If irritation develops, discontinue use'.
5205	WITHANIA SOMNIFERA	A, E, H	The medicine requires the following warning statement on the label:
			- (WITHANIA) 'If you are pregnant, or considering becoming pregnant, do not take without consulting a health professional' (or words to that effect)
			unless:
			(a) the plant part is root;
			(b) the plant preparation is an extract;
			(c) the extraction solvents are only water, ethanol or methanol; and
			(d) the maximum recommended daily dose of the medicine contains no more than the equivalent quantity of 12 g dry root.
5206	WOLFIPORIA COCOS	A, E, H	
5207	WOOL ALCOHOLS	Е	Only for use in topical medicines for

			dermal application.
5208	WOOL FAT	A, E	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing from time to time.

Volume 6

olume 6			
5209	XANTHAN GUM	Е	
5210	XANTHIUM SIBIRICUM	A, H	
5211	XANTHIUM STRUMARIUM	A, H	
5212	XANTHOMONA CAMPESTRIS	A, H	
5213	XEROPHYLLUM ASPHODELOIDES	A, H	
5214	XYLENE	E	The residual solvent limit for xylene is 21.7 mg per maximum recommended daily dose.
			The concentration in the medicine must be no more than 0.217%.
5215	XYLITOL	E	
5216	XYLOSE	Е	
5217	YAM	Е	
5218	YARROW HERB DRY	A, H	
5219	YARROW HERB POWDER	A, H	
5220	YEAST AUTOLYSATE	Е	
5221	YEAST DRIED	A, E, H	
5222	YELLOW 2G	Е	Permitted for use only as a colour for topical use.
5223	YELLOW BEESWAX	E	
5224	YELLOW MERCURIC OXIDE	Н	Only for use as an active homoeopathic ingredient.
5225	YELLOW SOFT PARAFFIN	Α, Ε	Only for use in topical medicines for dermal application.
			When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing from time to time.
5226	YLANG YLANG OIL	А, Е, Н	
5227	YUCCA BACCATA	A, H	
5228	YUCCA ELATA	A, H	
5229	YUCCA FILAMENTOSA	A, H	
5230	YUCCA GLORIOSA	A, H	

			volume (
5231	Z-BETA-DAMASCONE	E	Z – beta damascone must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation.
			The total concentration of flavour proprietary excipient formulations containing Z – beta damascone must not be more than 5% of the total medicine.
5232	ZANTHOXYLUM AMERICANUM	A, H	
5233	ZANTHOXYLUM BUNGEANUM	A, E, H	
5234	ZANTHOXYLUM CLAVA-HERCULIS	A, H	
5235	ZANTHOXYLUM NITIDUM	A, H	
5236	ZANTHOXYLUM PIPERITUM	A, H	
5237	ZANTHOXYLUM SIMULANS	A, H	
5238	ZEA MAYS	A, E, H	
5239	ZEAXANTHIN	A, E	
5240	ZEIN	Е	
5241	ZINC	Н	Only for use as an active homoeopathic ingredient.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, th medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect)'.
5242	ZINC AMINO ACID CHELATE	A, E, H	When used internally, zinc is a mandatory component of zinc amino aci chelate.
			The concentration of zinc in zinc amino acid chelate must be no more than 30%.

Therapeutic Goods (Permissible Ingredients) Determination (No. 2) 2024

Volume 6

Volume 6

			 When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: - (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5243	ZINC ASCORBATE	A, E, H	When used internally, zinc is a mandatory component of zinc ascorbate. When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: - (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5244	ZINC ASCORBATE MONOHYDRATE	А, Е, Н	When used internally, zinc is a mandatory component of zinc ascorbate monohydrate. When for internal use, the maximum recommended daily dose must be no
			 more than 50mg of zinc. When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: - (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if

			taken in large amounts or for a long
			period (or words to that effect)'.
5245	ZINC CHLORIDE	А, Е, Н	The concentration of zinc chloride in the medicine must be no more than 5%.
			When used internally, zinc is a mandatory component of zinc chloride.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5246	ZINC CITRATE	А, Е, Н	When used internally, zinc is a mandatory component of zinc citrate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5247	ZINC CITRATE DIHYDRATE	A, E, H	When used internally, zinc is a mandatory component of zinc citrate dihydrate.

Volume 6

Therapeutic Goods (Permissible Ingredients) Determination (No. 2) 2024

Volume 6

			 When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc. When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5248	ZINC CITRATE TRIHYDRATE	A, E, H	When used internally, zinc is a mandatory component of zinc citrate trihydrate. When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5249	ZINC DIASPARTATE	А	When used internally, zinc is a mandatory component of zinc diaspartate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or

			for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5250	ZINC GLUCONATE	А, Е, Н	When used internally, zinc is a mandatory component of zinc gluconate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5251	ZINC GLYCINATE	А	When used internally, zinc is a mandatory component of Zinc glycinate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5252	ZINC GLYCINATE MONOHYDRATE	A	When used internally, zinc is a mandatory component of Zinc glycinate monohydrate.

Volume 6

			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc. When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: - (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect)'.
5253	ZINC LACTATE	Ε	Only for use in topical and dental medicines and not to be included in medicines intended for use in the eye.
			The concentration of zinc lactate in a medicine intended for topical use should not be more than 2%.
			The concentration of zinc lactate in a medicine for 'dental' use in toothpaste medicines must not be more than 2.5%.
			Zinc lactate is not to be included in dental/toothpaste medicines intended for use by children less than 12 years old.
			Medicines containing zinc lactate for dental use require the following warning statement (or words to the same effect) on the medicine label:
			- (CHILD3) 'Use in children under 12 years is not recommended'.
5254	ZINC LACTATE DIHYDRATE	Е	Only for use in topical and dental medicines and not to be included in medicines intended for use in the eye.
			The concentration of zinc lactate dihydrate in a medicine intended for topical use should not be more than 2%.
			The concentration of zinc lactate dihydrate in a medicine for 'dental' use in toothpaste medicines must not be more than 2.5%.

Vol	lume	6

			Zinc lactate dihydrate is not to be included in dental/toothpaste medicines intended for use by children less than 12 years old.
			Medicines containing zinc lactate dihydrate for dental use require the following warning statement (or words to the same effect) on the medicine label:
			- (CHILD3) 'Use in children under 12 years is not recommended'.
5255	ZINC LYSINATE	А	When used internally, zinc is a mandatory component of Zinc lysinate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if
			taken in large amounts or for a long period (or words to that effect).'
5256	ZINC METHIONINE SULFATE	А	For topical use, the concentration of zinc methionine sulfate must be no more than 5%.
			When used internally, zinc is a mandatory component of zinc methionine sulfate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than
			25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or

Volume 6

			for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5257	ZINC MYRISTATE	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 0.1%.
5258	ZINC OXIDE	A, E, H	When used internally, zinc is a mandatory component of zinc oxide.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR
			-'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period' (or words to that effect).
			When used in primary sunscreen products, the following warning statements are required on the label:
			- (AVOID) 'Avoid prolonged exposure ir the sun' (or words to this effect); and
			- (SUNPRO) 'Wear protective clothing - hats and eyewear when exposed to the sun' (or words to this effect).
5259	ZINC PARA-PHENOLSULFONATE	E	Only permitted for use in topical medicines for dermal use.
			The concentration of zinc para- phenolsulfonate in the medicine must not exceed 5%.
5260	ZINC STEARATE	E	When used internally, zinc is a mandatory component of zinc stearate.

Volu	ıme 6
1 010	

			The percentage of zinc from zinc stearate should be calculated based on the molecular weight of zinc stearate. When for internal use, the maximum recommended daily dose must not provide more than 50 milligrams of zinc.
			When for internal use and the maximum recommended daily dose is more than 25 milligrams but not more than 50 milligrams of zinc, the medicine requires the following warning statement (or words to the same effect) on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period'; or
		'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period'.	
5261	ZINC SUCCINATE	A, E, H	When used internally, zinc is a mandatory component of zinc succinate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' or
			- 'WARNING: Contains zinc which may be dangerous if taken in large amounts on for a long period (or words to that effect).'
5262	ZINC SULFATE	Α, Ε	For topical use, the concentration of zinc sulfate must be no more than 5%.
			For internal use, zinc is a mandatory component of zinc sulfate.

Volume 6

oranne o			
			 When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc. When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5263	ZINC SULFATE HEPTAHYDRATE	Α, Ε	For topical use, the concentration of zinc sulfate must be no more than 5%. For internal use, zinc is a mandatory component of zinc sulfate heptahydrate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR
			- 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5264	ZINC SULFATE HEXAHYDRATE	A, E, H	For topical use, the concentration of zinc sulfate must be no more than 5%.
			For internal use, zinc is a mandatory component of zinc sulfate hexahydrate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.

Vol	lume	6
v U	luine	v

			v ordinie (
			 When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label: - (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR
			- 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5265	ZINC SULFATE MONOHYDRATE	A, E, H	When the route of administration is topical the concentration of zinc sulfate in the medicine must be no more than 5%.
			When the medicine is for internal use, zinc is a mandatory component of zinc sulfate monohydrate.
			When for internal use, the maximum recommended daily dose must be no more than 50mg of zinc.
			When for internal use and the maximum recommended daily dose is more than 25mg but no more than 50mg of zinc, the medicine requires the following warning statement on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period.' OR
			- 'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period (or words to that effect).'
5266	ZINC VALERATE	Н	Only for use as an active homoeopathic ingredient.
			When for internal use, zinc is a mandatory component of zinc valerate.
			The percentage of zinc from zinc valerate should be calculated based on the molecular weight of zinc valerate.

Volume	6	

			When for internal use, the maximum recommended daily dose must not provide more than 50 milligrams of zinc.
			When for internal use and the maximum recommended daily dose is more than 25 milligrams but not more than 50 milligrams of zinc, the medicine requires the following warning statement (or words to the same effect) on the medicine label:
			- (ZINC) 'WARNING: May be dangerous if taken in large amounts or for a long period'; or
			'WARNING: Contains zinc which may be dangerous if taken in large amounts or for a long period'.
5267	ZINGERONE	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
5268	ZINGIBER OFFICINALE	A, E, H	When for oral use AND the extract ratio is equal to or more than 25:1 AND the equivalent dry weight per dosage unit is equal to or more than 2g, the medicine requires the following warning statement on the medicine label:
			- (GINGER) 'Individuals taking anticoagulants should seek medical advice before taking this medicine.' AND 'Individuals at risk of bleeding problems should seek advice from their healthcare practitioner prior to taking this medicine'.
5269	ZIZIPHUS JUJUBA	A, H	
5270	ZIZIPHUS JUJUBA VAR. SPINOSA	A, H	
5271	ZIZYPHUS SATIVA	A, H	
5272	ZOSTERA MARINA	A, H	